

PRO-105.0  
6868/75530

-1-

## EXOGENOUS NUCLEIC ACID DETECTION

### Description

#### Cross-Reference to Related Applications

5                   This application is a continuation-in-part  
of U.S. Serial No. 09/358,972, filed on July 21,  
1999, which is a continuation-in-part of U.S. Serial  
No. 09/252,436, filed on February 18, 1999, which is  
a continuation-in-part of U.S. Serial No. 09/042,287,  
10       filed March 13, 1998, all of which are incorporated  
herein by reference.

#### Field of the Invention

                  The invention relates to nucleic acid  
15       detection. More specifically, the invention relates  
to the detection of a predetermined exogenous nucleic  
acid target sequence in a nucleic acid target/probe  
hybrid, and the various applications of such  
detection.

20

#### Background of the Invention

                  Methods to detect nucleic acids provide a  
foundation upon which the large and rapidly growing  
field of molecular biology is built. There is  
25       widespread application of such general methods to the  
detection of specific, exogenous nucleic acids.  
There is constant need for alternative methods and  
products. The reasons for selecting one method over  
another are varied, and include a desire to avoid  
30       radioactive materials, the lack of a license to use a

technique, the cost or availability of reagents or equipment, the desire to minimize the time spent or the number of steps, the accuracy or sensitivity needed for a certain application, the ease of  
5 analysis, or the ability to automate the process.

The detection of nucleic acids, including specific exogenous nucleic acids, is often a portion of a process rather than an end in itself. There are many applications of the detection of nucleic acids  
10 in the art, and new applications are always being developed. The ability to detect and quantify exogenous nucleic acids is useful in detecting microorganisms and viruses and biological molecules (e.g. non-native promoter or terminator sequences or  
15 foreign genes) in a biological sample, and thus affects many fields, including human and veterinary medicine, food processing and environmental testing. Additionally, the detection and/or quantification of specific biomolecules from biological samples (e.g.  
20 tissue, sputum, urine, blood, semen, saliva) has applications in medicine and forensic science.

Hybridization methods to detect nucleic acids are dependent upon knowledge of the nucleic acid sequence. Many known nucleic acid detection  
25 techniques depend upon specific nucleic acid hybridization in which an oligonucleotide probe is hybridized or annealed to nucleic acid in the sample or on a blot, and the hybridized probes are detected.

A traditional type of process for the  
30 detection of hybridized nucleic acid uses labeled nucleic acid probes to hybridize to a nucleic acid

sample. For example, in a Southern blot technique, a nucleic acid sample is separated in an agarose gel based on size and affixed to a membrane, denatured, and exposed to a labeled nucleic acid probe under hybridizing conditions. If the labeled nucleic acid probe forms a hybrid with the nucleic acid on the blot, the label is bound to the membrane. Probes used in Southern blots have been labeled with radioactivity, fluorescent dyes, digoxigenin, horseradish peroxidase, alkaline phosphatase and acridinium esters.

Another type of process for the detection of hybridized nucleic acid takes advantage of the polymerase chain reaction (PCR). The PCR process is well known in the art (U.S. Patent Nos. 4,683,195, 4,683,202, and 4,800,159). To briefly summarize PCR, nucleic acid primers, complementary to opposite strands of a nucleic acid amplification target sequence, are permitted to anneal to the denatured sample. A DNA polymerase (typically heat stable) extends the DNA duplex from the hybridized primer. The process is repeated to amplify the nucleic acid target. If the nucleic acid primers do not hybridize to the sample, then there is no corresponding amplified PCR product. In this case, the PCR primer acts as a hybridization probe. PCR-based methods are of limited use for the detection of nucleic acid of unknown sequence.

In a PCR method, the amplified nucleic acid product may be detected in a number of ways, e.g. incorporation of a labeled nucleotide into the

amplified strand by using labeled primers. Primers used in PCR have been labeled with radioactivity, fluorescent dyes, digoxigenin, horseradish peroxidase, alkaline phosphatase, acridinium esters, biotin and jack bean urease. PCR products made with unlabeled primers may be detected in other ways, such as electrophoretic gel separation followed by dye-based visualization.

Enzymes having template-specific polymerase activity for which some 3'→5' depolymerization activity has been reported include *E. coli* DNA Polymerase (Deutscher and Kornberg, *J. Biol. Chem.*, **244**(11):3019-28 (1969)), T7 DNA Polymerase (Wong et al., *Biochemistry* **30**:526-37 (1991); Tabor and Richardson, *J. Biol. Chem.* **265**: 8322-28 (1990)), *E. coli* RNA polymerase (Rozovskaya et al., *Biochem. J.* **224**:645-50 (1994)), AMV and RLV reverse transcriptases (Srivastava and Modak, *J. Biol. Chem.* **255**: 2000-4 (1980)), and HIV reverse transcriptase (Zinnen et al., *J. Biol. Chem.* **269**:24195-202 (1994)). A template-dependent polymerase for which 3' to 5' exonuclease activity has been reported on a mismatched end of a DNA hybrid is phage 29 DNA polymerase (de Vega, M. et al. *EMBO J.*, **15**:1182-1192, 1996)

There is a need for highly sensitive, diagnostic applications that are capable of determining the number of virus molecules present in a body ("viral load"). For example, the presence of viral particles in the circulation system or in specific tissues is a means of monitoring the

severity of viral infection. Several methods are currently used in the art for determining viral load. U.S. Patent No. 5,667,964 discloses a method for the determination of the number of HIV-1 infected patient cells using reactive oxygen-intermediate generators. U.S. Patent No. 5,389,512 discloses a method for determining the relative amount of a viral nucleic acid segment in a sample using PCR.

G. Garinis et al., *J. Clin. Lab. Anal.* 13:122-5 (1999) compare the determination of viral load results using an enzyme-linked immunosorbent assay (ELISA), a recombinant immunoblot assay (RIBA), and a reverse transcriptase polymerase chain reaction method (RT-PCR) in the detection of hepatitis C virus (HCV) infection in haemodialysis patients. The quantitative hepatitis HCV RT-PCR assay had a detection level of about 2,000 viral copies/mL serum. Holguin et al., *Eur. J. Clin. Microbiol. Infect. Dis.* 18:256-9 (1999) compare plasma HIV-1 RNA levels using several commercially available assays, namely the second-generation HIV-1 branched DNA assay, the Nuclisens assay, the Amplicor® Monitor reverse transcriptase polymerase chain reaction assay, and the Ultradirect Monitor. Differing values were noted in comparing results among these various assays. Boriskin et al., *Arch. Dis. Child.* 80:132-6 (1999) used a nested polymerase chain reaction to measure HIV-1 proviral DNA and CMV genomic DNA in peripheral blood leukocytes of children infected with HIV-1. There remains a need for a reliable means to detect and quantify viral load. There is a demand for

methods to determine viral load when the quantities of viral particles are very low.

There is a need for alternative methods for detection of nucleic acid hybrids. There is a demand  
5 for highly sensitive methods that are useful for determining the presence or absence of specific nucleic acid sequences that are non-native or "exogenous" to an organism's nucleic acid. For example, there is a need to determine the presence of  
10 non-native nucleic acid present in a cell, both when the non-native nucleic acid is incorporated into the native nucleic acid and when it is not incorporated. For example, there is a need for methods to determine viral load that are able to reliably detect as few as  
15 10 copies of a virus present in a body, tissue, fluid, or other biological sample. There is great demand for methods to determine the presence of a mutant virus, e.g. a drug-resistant mutant, in a biological sample containing a viral population.  
20 There is great demand for methods to determine the presence or absence of non-native sequences unique to a particular species in a sample, for example the identification of bacterial contamination present in a primarily non-bacterial biological sample. There  
25 is also great demand for methods that are more highly sensitive than the known methods, methods that are highly reproducible and automatable.

It would be beneficial if another method were available for detecting the presence of a  
30 sought-after, predetermined exogenous target nucleotide sequence. It would also be beneficial if

such a method were operable using a sample size of the microgram to picogram scale. It would further be beneficial if such a detection method were capable of providing multiple analyses in a single assay (multiplex assays). The disclosure that follows provides one such method.

#### Brief Summary of the Invention

A method of this invention is used to determine the presence or absence of a predetermined exogenous nucleic acid target sequence in a nucleic acid sample. Such a method utilizes an enzyme that depolymerizes the 3'-terminus of an oligonucleotide probe hybridized to a nucleic acid target sequence to release one or more identifier nucleotides whose presence can then be determined.

One embodiment of the invention contemplates a method for determining the presence or absence of a predetermined, exogenous nucleic acid target sequence in a nucleic acid sample. More than one such predetermined target sequence can also be present in the sample being assayed, and the presence or absence of more than one predetermined nucleic acid target sequence can be determined. The embodiment comprises the following steps.

A treated sample is provided that may contain a predetermined nucleic acid target sequence hybridized with a nucleic acid probe that includes an identifier nucleotide in the 3'-terminal region. The treated sample is admixed with a depolymerizing amount of an enzyme whose activity is to release one

or more nucleotides from the 3'-terminus of a hybridized nucleic acid probe to form a treated reaction mixture. The type of enzyme used for nucleotide release will be further identified and described herein. The treated reaction mixture is maintained under depolymerizing conditions for a time period sufficient to permit the enzyme to depolymerize hybridized nucleic acid and release identifier nucleotides therefrom. The presence of released identifier nucleotides is analyzed to obtain an analytical output, the analytical output indicating the presence or absence of the nucleic acid target sequence. The analytical output is obtained by various techniques as discussed herein.

It is contemplated that an analytical output of a method of the invention can be obtained in a variety of ways. The analytical output can be ascertained by luminescence spectroscopy. In some preferred embodiments, analysis for released 3'-terminal region indicator nucleotides comprises the detection of ATP, either by a luciferase detection system (luminescence spectroscopy) or an NADH detection system (absorbance or fluorescence spectroscopy). In particularly preferred embodiments, ATP molecules are formed by a phosphate transferring step, for example using an enzyme such as NDPK in the presence of ADP, with the phosphate group originating from the nucleotide triphosphates produced by the depolymerizing step. In other embodiments, the ATP produced is amplified to form a plurality of ATP molecules. In the ATP detection



embodiments, the enzyme nucleoside diphosphate kinase (NDPK) is typically present in the depolymerization reaction and functions to convert released nucleotides and added ADP into ATP, and thus  
5 reactions where the two enzymes are present together, are denoted as a "one pot" method.

In an alternative embodiment, the analytical output is obtained by fluorescence spectroscopy. Fluorescence can be incorporated or  
10 added to a probe in a number of ways known in the art. For example, it is contemplated that an identifier nucleotide includes a fluorescent label. An identifier nucleotide can be fluorescently labeled prior to, or after, release of the identifier  
15 nucleotide. It is also contemplated that other than a released identifier nucleotide contains a fluorescent tag. In such an embodiment, the release of nucleotides in a process of the invention is ascertained by a determination of a difference in the  
20 length of the polynucleotide probe, for example by capillary electrophoresis imaged by a fluorescent tag at the 5' terminus of the probe or in a region other than the 3' terminal region.

In an alternative embodiment the analytical  
25 output is obtained by mass spectrometry. It is preferred here that an identifier nucleotide be a nucleotide analog or a labeled nucleotide and have a molecular mass that is different from the mass of a usual form of that nucleotide, although a difference  
30 in mass is not required. It is also noted that with a fluorescent-labeled identifier nucleotide, the

analytical output can also be obtained by mass spectrometry. It is also contemplated that the analysis of released nucleotide be conducted by ascertaining the difference in mass of the probe  
5 after a depolymerization step of a process of the invention.

In another alternative embodiment, the analytical output is obtained by absorbance spectroscopy. Such analysis monitors the absorbance  
10 of light in the ultraviolet and visible regions of the spectrum to determine the presence of absorbing species. In one aspect of such a process, released nucleotides are separated from hybridized nucleic acid and other polynucleotides by chromatography  
15 (e.g. HPLC or GC) or electrophoresis (e.g. PAGE or capillary electrophoresis). Either the released identifier nucleotide or the remainder of the probe can be analyzed to ascertain the release of the identifier nucleotide in a process of the invention.  
20 In another aspect of such a process a label may be incorporated in the analyzed nucleic acid.

In a contemplated embodiment, a sample to be assayed for the presence or absence of an exogenous nucleic acid target sequence is admixed  
25 with one or more nucleic acid probes under hybridizing conditions to form a hybridization composition. The 3'-terminal region of the nucleic acid probe hybridizes with partial or total complementarity to the exogenous nucleic acid target  
30 sequence when that sequence is present in the sample. The 3'-terminal region of the nucleic acid probe

includes an identifier nucleotide. The hybridization composition is maintained under hybridizing conditions for a time period sufficient to form a treated sample that may contain said predetermined nucleic acid target sequence hybridized with a nucleic acid probe. The treated sample is admixed with a depolymerizing amount of an enzyme whose activity is to release one or more nucleotides from the 3'-terminus of a hybridized nucleic acid probe to form a treated reaction mixture. The treated reaction mixture is maintained under depolymerizing conditions for a time period sufficient to permit the enzyme to depolymerize hybridized nucleic acid and release identifier nucleotides therefrom. The presence of released identifier nucleotides is analyzed to obtain an analytical output, the analytical output indicating the presence or absence of the nucleic acid target sequence. The analytical output may be obtained by various techniques as discussed above.

One method of the invention contemplates interrogating the presence or absence of a specific base in an exogenous nucleic acid target sequence in a sample to be assayed, and comprises the following steps.

A hybridization composition is formed by admixing a sample to be assayed with one or more nucleic acid probes under hybridizing conditions. The sample to be assayed may contain an exogenous nucleic acid target sequence to be interrogated. The nucleic acid target comprises at least one base whose

presence or absence is to be identified. The hybridization composition includes at least one nucleic acid probe that is substantially complementary to the nucleic acid target sequence and comprises at least one predetermined nucleotide at an interrogation position, and an identifier nucleotide in the 3' terminal region.

A treated sample is formed by maintaining the hybridization composition under hybridizing conditions for a time period sufficient for base pairing to occur when a probe nucleotide at an interrogation position is aligned with a base to be identified in the target sequence. A treated reaction mixture is formed by admixing the treated sample with an enzyme whose activity is to release one or more identifier nucleotides from the 3' terminus of a hybridized nucleic acid probe to depolymerize the hybrid. The treated reaction mixture is maintained under depolymerizing conditions for a time period sufficient to permit the enzyme to depolymerize the hybridized nucleic acid and release an identifier nucleotide.

An analytical output is obtained by analyzing for the presence or absence of released identifier nucleotides. The analytical output indicates the presence or absence of the specific base or bases to be identified. The analytical output is obtained by various techniques as discussed herein. Preferably, an identifier nucleotide is at the interrogation position.

In one aspect of a method of the invention, the nucleic acid target sequence is selected from the group consisting of deoxyribonucleic acid and ribonucleic acid.

5 In another aspect of the invention, the sample containing a plurality of target nucleic acid sequences is admixed with a plurality of the nucleic acid probes. Several analytical outputs can be obtained from such multiplexed assays.

10 In a first embodiment, the analytical output obtained when at least one nucleic acid probe hybridizes with partial complementarity to one target nucleic acid sequence is greater than the analytical output when all of the nucleic acid probes hybridize  
15 with total complementarity to their respective nucleic acid target sequences. In a second embodiment, the analytical output obtained when at least one nucleic acid probe hybridizes with partial complementarity to one target nucleic acid sequence  
20 is less than the analytical output when all of the nucleic acid probes hybridize with total complementarity to their respective nucleic acid target sequences. In a third embodiment, the analytical output obtained when at least one nucleic  
25 acid probe hybridizes with total complementarity to one nucleic acid target sequence is greater than the analytical output when all of the nucleic acid probes hybridize with partial complementarity to their respective nucleic acid target sequences. In a  
30 fourth embodiment, the analytical output obtained when at least one nucleic acid probe hybridizes with

total complementarity to one target nucleic acid  
sequence is less than the analytical output when all  
of the nucleic acid probes hybridize with partial  
complementarity to their respective nucleic acid  
5 target sequences. The depolymerizing enzymes are as  
described herein.

Yet another embodiment of the invention  
contemplates a method for determining the presence or  
absence of a first exogenous nucleic acid target in a  
10 nucleic acid sample that may contain that target or  
may contain a substantially identical second target.  
For example, the second target may have a base  
substitution, deletion or addition relative to the  
first nucleic acid target. This embodiment comprises  
15 the following steps.

A sample to be assayed is admixed with one  
or more nucleic acid probes under hybridizing  
conditions to form a hybridization composition. The  
first and second nucleic acid targets each comprise a  
20 region of sequence identity except for at least a  
single nucleotide at a predetermined position that  
differs between the targets. The nucleic acid probe  
is substantially complementary to the nucleic acid  
target region of sequence identity and comprises at  
25 least one nucleotide at an interrogation position.  
An interrogation position of the probe is aligned  
with the predetermined position of a target when a  
target and probe are hybridized. The probe also  
includes an identifier nucleotide in the 3'-terminal  
30 region.

The hybridization composition is maintained under hybridizing conditions for a time period sufficient to form a treated sample wherein the nucleotide at the interrogation position of the probe  
5 is aligned with the nucleotide at the predetermined position in the region of identity of the target.

A treated reaction mixture is formed by admixing the treated sample with a depolymerizing amount of an enzyme whose activity is to release one  
10 or more nucleotides from the 3'-terminus of a hybridized nucleic acid probe. The reaction mixture is maintained under depolymerization conditions for a time period sufficient to permit the enzyme to depolymerize the hybridized nucleic acid and release  
15 the identifier nucleotide.

An analytical output is obtained by analyzing for the presence or absence of released identifier nucleotides. The analytical output indicates the presence or absence of the nucleotide  
20 at the predetermined region, and; thereby, the presence or absence of a first nucleic acid target.

One aspect of the above method is comprised of a first probe and a second probe. The first probe comprises a nucleotide at an interrogation position  
25 that is complementary to a first nucleic acid target at a predetermined position. The second probe comprises a nucleotide at an interrogation position that is complementary to a second nucleic acid target at a predetermined position.

30 In another aspect of a process of the invention, the depolymerizing enzyme, whose activity

is to release nucleotides, is a template-dependent polymerase, whose activity is to depolymerize hybridized nucleic acid, whose 3'-terminal nucleotide is matched, in the 3'→5' direction in the presence of pyrophosphate ions to release one or more nucleotides. Thus, the enzyme's activity is to depolymerize hybridized nucleic acid to release nucleotides under depolymerizing conditions. Preferably, this enzyme depolymerizes hybridized nucleic acids whose bases in the 3'-terminal region of the probe are matched with total complementarity to the corresponding bases of the nucleic acid target.

In an alternative aspect of the process of the invention, the depolymerizing enzyme, whose activity is to release nucleotides, exhibits a 3'→5' exonuclease activity in which hybridized nucleic acids having one or more mismatched bases at the 3'-terminus of the hybridized probe are depolymerized. Thus, the enzyme's activity is to depolymerize hybridized nucleic acid to release nucleotides under depolymerizing conditions. In this embodiment, the hybrid can be separated from the free probe prior to enzyme treatment. In some embodiments, an excess of target may be used so that the concentration of free probe in the enzyme reaction is extremely low.

In still another alternative aspect of a process of the invention, the depolymerizing enzyme exhibits a 3' to 5' exonuclease activity on a double-stranded DNA substrate having one or more matched bases at the 3' terminus of the hybrid. The enzyme's



activity is to depolymerize hybridized nucleic acid to release nucleotides containing a 5' phosphate under depolymerizing conditions.

In particularly preferred embodiments, ATP molecules are formed by a phosphate transferring step, (e.g. using the enzyme NDPK in the presence of ADP), from the deoxynucleoside triphosphates (dNTPs) produced by the depolymerizing step. In some embodiments, the ATP can be amplified to form a plurality of ATP molecules. Thermostable nucleoside diphosphate kinases are particularly preferred when an NDPK enzyme is used.

In one aspect of the invention, the nucleic acid sample to be assayed is obtained from a biological sample that is a solid or liquid.

In one aspect of the method, the predetermined nucleic acid target sequence is present in the sample for the purpose of gene therapy.

In one aspect of the method, the predetermined nucleic acid target sequence is a microbial or viral nucleic acid.

In some preferred embodiments of the invention, the predetermined nucleic acid target sequence is a viral nucleic acid. Viral load, the amount of virus present, can be determined from the magnitude of the analytical output from a predetermined amount of biological sample such as animal fluid or tissue.

In some preferred embodiments, the presence or absence of a mutation in the viral genome can be determined.

In another aspect of the method, the nucleic acid sample is obtained from a food source. In one process of the method, the food source is a plant or is derived from plant material, and the predetermined nucleic acid target sequence is a sequence not native to that plant. In one aspect of the method, the nucleic acid sequence not native to the subject plant is a transcription control sequence. In one preferred embodiment of the invention, the transcription control sequence is the 35S promoter or the NOS terminator, or both.

A still further embodiment of the invention contemplates determining the presence or absence of an exogenous nucleic acid target sequence in a nucleic acid sample with a probe that is hybridized to the target and then modified to be able to form a hairpin structure. This embodiment comprises the following steps.

A treated sample is provided that contains a nucleic acid sample that may include an exogenous nucleic acid target sequence having an interrogation position hybridized with a nucleic acid probe. The probe is comprised of at least two sections. The first section contains the probe 3'-terminal about 10 to about 30 nucleotides. These nucleotides are complementary to the target strand sequence at positions beginning about 1 to about 30 nucleotides downstream of the interrogation position. The second section of the probe is located at the 5'-terminal region of the probe and contains about 10 to about 20 nucleotides of the target sequence. This sequence

spans the region in the target from the nucleotide at  
or just upstream (5') of the interrogation position,  
to the nucleotide just upstream to where the 3'-  
terminal nucleotide of the probe anneals to the  
5 target. An optional third section of the probe, from  
zero to about 50 nucleotides, and preferably about  
zero to about 20 nucleotides in length, and  
comprising a sequence that does not hybridize with  
either the first or second section, is located  
10 between the first and second sections of the probe.

The probe of the treated sample is extended  
in a template-dependent manner, as by admixture with  
dNTPs and a template-dependent polymerase, at least  
through the interrogation position, thereby forming  
15 an extended probe/target hybrid. In a preferred  
embodiment, the length of the probe extension is  
limited by omission from the extension reaction of a  
dNTP complementary to a nucleotide of the target  
sequence that is present upstream of the  
20 interrogation position and absent between the  
nucleotide complementary to the 3'-end of the  
interrogation position.

The extended probe/target hybrid is  
separated from any unreacted dNTPs. The extended  
25 probe/target hybrid is denatured to separate the  
strands. The extended probe strand is permitted to  
form a hairpin structure.

A treated reaction mixture is formed by  
admixing the hairpin structure-containing composition  
30 with a depolymerizing amount of an enzyme whose  
activity is to release one or more nucleotides from

the 3'-terminus of an extended probe hairpin structure. The reaction mixture is maintained under depolymerizing conditions for a time period sufficient for the depolymerizing enzyme to release 3'-terminus nucleotides, and then analyzed for the presence of released identifier nucleotides. The analytical output indicates the presence or absence of the exogenous nucleic acid target sequence.

A still further embodiment of the invention, termed REAPER™, also utilizes hairpin structures. This method contemplates determining the presence or absence of an exogenous nucleic acid target sequence, or a specific base within the target sequence, in a nucleic acid sample, and comprises the following steps. A treated sample is provided that contains a nucleic acid sample that may include an exogenous nucleic acid target sequence hybridized with a first nucleic acid probe strand.

The hybrid is termed the first hybrid. The first probe is comprised of at least two sections. The first section contains the probe 3'-terminal about 10 to about 30 nucleotides that are complementary to the target nucleic acid sequence at a position beginning about 5 to about 30 nucleotides downstream of the target interrogation position. The second section of the first probe contains about 5 to about 30 nucleotides that are a repeat of the target sequence from the interrogation position to about 10 to about 30 nucleotides downstream of the interrogation position, and does not hybridize to the first section of the probe. An optional third

section of the probe, located between the first and second sections of the probe, is zero to about 50 nucleotides, preferably up to about 20 nucleotides, in length and comprises a sequence that does not hybridize to either the first or second section.

The first hybrid in the treated sample is extended at the 3'-end of the first probe, thereby extending the first probe past the interrogation position and forming an extended first hybrid whose sequence includes an interrogation position. The extended first hybrid is comprised of the original target nucleic acid and extended first probe. The extended first hybrid is then denatured in an aqueous composition to separate the two nucleic acid strands of the hybridized duplex and form an aqueous solution containing a separated target nucleic acid and a separated extended first probe.

A second probe, that is about 10 to about 2000 nucleotides, preferably about 10 to about 200 nucleotides, most preferably about 10 to about 30 nucleotides, in length and is complementary to the extended first probe at a position beginning about 5 to about 2000, preferably about 5 to about 200, nucleotides downstream of the interrogation position in extended first probe, is annealed to the extended first probe, thereby forming the second hybrid. The second hybrid is extended at the 3'-end of the second probe until that extension reaches the 5'-end of the extended first probe, thereby forming a second extended hybrid whose 3'-region includes an identifier nucleotide. In preferred embodiments the

extending polymerase for both extensions does not add a nucleotide to the 3' end that does not have a corresponding complementary nucleotide in the template.

5           An aqueous composition of the extended second hybrid is denatured to separate the two nucleic acid strands. The aqueous composition so formed is cooled to form a "hairpin structure" from the separated extended second probe when the target  
10           sequence is present in the original nucleic acid sample.

          A treated reaction mixture is formed by admixing the hairpin structure-containing composition with a depolymerizing amount of an enzyme whose  
15           activity is to release one or more nucleotides from the 3'-terminus of a nucleic acid hybrid. The reaction mixture is maintained under depolymerizing conditions for a time period sufficient to release 3'-terminal region identifier nucleotides, and then  
20           analyzed for the presence of released identifier nucleotides. The analytical output indicates the presence or absence of the exogenous nucleic acid target sequence.

          The present invention has many benefits and  
25           advantages, several of which are listed below.

          One benefit of the invention is that, in some embodiments, nucleic acid hybrids can be detected with very high levels of sensitivity without the need for radiochemicals or electrophoresis.

30           An advantage of the invention is that the presence or absence of one or more exogenous target

nucleic acid(s) can be detected reliably,  
reproducibly, and with great sensitivity.

5 A further benefit of the invention is that  
quantitative information can be obtained about the  
amount of exogenous target nucleic acid sequence in a  
sample and a large variety of sample types can be  
used.

10 A further advantage of the invention is  
that very slight differences in exogenous nucleic  
acid sequence are detectable, including single  
nucleotide polymorphisms (SNPs).

15 Yet another benefit of the invention is  
that the presence or absence of a number of exogenous  
target nucleic acid sequences can be determined in  
the same assay.

Yet another advantage of the invention is  
that the presence or absence of an exogenous target  
nucleic acid can be determined with a small number of  
reagents and manipulations.

20 Another benefit of the invention is that  
the processes lend themselves to automation.

25 Still another benefit of the invention is  
its flexibility of use in many different types of  
applications and assays including, but not limited  
to, determination of viral load, determination of  
viral type, species identification, sample  
contamination, and analysis of forensic samples.

30 Still further benefits and advantages of  
the invention will become apparent to the skilled  
worker from the disclosure that follows.

Brief Description of the Figures

In the drawings forming a portion of this disclosure,

5           Fig. 1 illustrates the annealing of the 10865 oligonucleotide (SEQ ID NO:76) to 10870 wild type (SEQ ID NO:77) and 10994 mutant (SEQ ID NO:78) oligonucleotides utilized in rolling circle amplification as Fig. 1A and Fig. 1B, respectively.  
10       Also shown are the annealing (hybridization) of oligonucleotide 10866 (SEQ ID NO:81) to oligonucleotide 10865, as well as the hybridization of oligonucleotide probe 10869 (SEQ ID NO:79) to oligonucleotide 10870 and of oligonucleotide probe  
15       10989 (SEQ ID NO:80) to oligonucleotide 10994 as representations of the binding of those probes to the respective amplified sequences. Arcuate lines in oligonucleotide 10865 are used to help illustrate the shape that oligonucleotide 10865 can assume when  
20       hybridized with either of oligonucleotides 10870 or 10994.

          Fig. 2. illustrates the Reaper™ assay as described in Example 21. Fig. 2A illustrates the first hybrid formed by the annealing of nucleic acid  
25       target SEQ ID NO:67 (286) to first probe SEQ ID NO:68 (287). An arrow points to an interrogation position in 286.

          Fig. 2B illustrates the first extended hybrid formed by the annealing of 286 to the extended  
30       287. Extended 287 is first extended probe SEQ ID NO:69 (288).



Fig. 2C illustrates the second hybrid formed by annealing of 288 from the denatured nucleic acid molecule shown in Fig. 2B to the second probe denoted SEQ ID NO:70 (289). An arrow points to the  
5 interrogation position in 288.

Fig. 2D illustrates the extended second hybrid formed by the annealing of 288 and the extended 289 strand denoted SEQ ID NO:71 (290).

Fig. 2E illustrates the 290 strand  
10 denatured from Fig. 2D and forming a hairpin structure. An arrow points to the interrogation position at the 3'-terminus of the hybrid.

#### 15 Definitions

To facilitate understanding of the invention, a number of terms are defined below.

"Nucleoside", as used herein, refers to a compound consisting of a purine [guanine (G) or  
20 adenine (A)] or pyrimidine [thymine (T), uridine (U) or cytidine (C)] base covalently linked to a pentose, whereas "nucleotide" refers to a nucleoside phosphorylated at one of its pentose hydroxyl groups. "XTP", "XDP" and "XMP" are generic designations for  
25 ribonucleotides and deoxyribonucleotides, wherein the "TP" stands for triphosphate, "DP" stands for diphosphate, and "MP" stands for monophosphate, in conformity with standard usage in the art. Subgeneric designations for ribonucleotides are  
30 "NMP", "NDP" or "NTP", and subgeneric designations for deoxyribonucleotides are "dNMP", "dNDP" or

"dNTP". Also included as "nucleoside", as used herein, are materials that are commonly used as substitutes for the nucleosides above such as modified forms of these bases (e.g. methyl guanine) or synthetic materials well known in such uses in the art, such as inosine.

A "nucleic acid," as used herein, is a covalently linked sequence of nucleotides in which the 3' position of the pentose of one nucleotide is joined by a phosphodiester group to the 5' position of the pentose of the next, and in which the nucleotide residues (bases) are linked in specific sequence; i.e., a linear order of nucleotides. A "polynucleotide," as used herein, is a nucleic acid containing a sequence that is greater than about 100 nucleotides in length. An "oligonucleotide," as used herein, is a short polynucleotide or a portion of a polynucleotide. An oligonucleotide typically contains a sequence of about two to about one hundred bases. The word "oligo" is sometimes used in place of the word "oligonucleotide".

A base "position" as used herein refers to the location of a given base or nucleotide residue within a nucleic acid.

A "nucleic acid of interest," as used herein, is any particular nucleic acid one desires to study in a sample.

The term "isolated" when used in relation to a nucleic acid or protein, refers to a nucleic acid sequence or protein that is identified and separated from at least one contaminant (nucleic acid

or protein, respectively) with which it is ordinarily associated in its natural source. Isolated nucleic acid or protein is present in a form or setting that is different from that in which it is found in  
5 nature. In contrast, non-isolated nucleic acids or proteins are found in the state they exist in nature.

As used herein, the term "purified" or "to purify" means the result of any process which removes some contaminants from the component of interest,  
10 such as a protein or nucleic acid. The percent of a purified component is thereby increased in the sample.

The term "wild-type," as used herein, refers to a gene or gene product that has the  
15 characteristics of that gene or gene product that is most frequently observed in a population and is thus arbitrarily designated the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" as used herein, refers to a gene or gene  
20 product that displays modifications in sequence and/or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product.

Nucleic acids are known to contain  
25 different types of mutations. As used herein, a "point" mutation refers to an alteration in the sequence of a nucleotide at a single base position.

A "single nucleotide polymorphism" or SNP, as used herein, is a variation from the most  
30 frequently occurring base at a particular nucleic acid position.

As used herein, the term "exogenous" with reference to a nucleic acid sequence is a nucleic acid that is not native in a sample. For example, a gene that was inserted into a cell is exogenous, or a virus that is present in a host cell is exogenous. It may be exogenous whether or not it is incorporated into the DNA of the host cell.

DNA molecules are said to have a "5'-terminus" (5' end) and a "3'-terminus" (3' end) because nucleic acid phosphodiester linkages occur to the 5' carbon and 3' carbon of the pentose ring of the substituent mononucleotides. The end of a polynucleotide at which a new linkage would be to a 5' carbon is its 5' terminal nucleotide. The end of a polynucleotide at which a new linkage would be to a 3' carbon is its 3' terminal nucleotide. A terminal nucleotide, as used herein, is the nucleotide at the end position of the 3'- or 5'-terminus. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide or polynucleotide, also can be said to have 5'- and 3'- ends. For example, a gene sequence located within a larger chromosome sequence can still be said to have a 5'- and 3'-end.

As used herein, the 3'-terminal region of the nucleic acid probe refers to the region of the probe including nucleotides within about 10 residues from the 3'-terminal position.

In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or "5'" relative to an element if they are bonded or would be bonded to the 5'-end of that

element. Similarly, discrete elements are  
"downstream" or "3'" relative to an element if they  
are or would be bonded to the 3'-end of that element.  
Transcription proceeds in a 5' to 3' manner along the  
5 DNA strand. This means that RNA is made by the  
sequential addition of ribonucleotide-5'-  
triphosphates to the 3'-terminus of the growing chain  
(with the elimination of pyrophosphate).

As used herein, the term "target nucleic  
10 acid" or "nucleic acid target" refers to a particular  
nucleic acid sequence of interest. Thus, the  
"target" can exist in the presence of other nucleic  
acid molecules or within a larger nucleic acid  
molecule.

As used herein, the term "nucleic acid  
15 probe" refers to an oligonucleotide or polynucleotide  
that is capable of hybridizing to another nucleic  
acid of interest. A nucleic acid probe may occur  
naturally as in a purified restriction digest or be  
20 produced synthetically, recombinantly or by PCR  
amplification. As used herein, the term "nucleic  
acid probe" refers to the oligonucleotide or  
polynucleotide used in a method of the present  
invention. That same oligonucleotide could also be  
25 used, for example, in a PCR method as a primer for  
polymerization, but as used herein, that  
oligonucleotide would then be referred to as a  
"primer". Herein, oligonucleotides or polynucleotides  
may contain a modified linkage such as a  
30 phosphorothioate bond.

As used herein, the terms "complementary" or "complementarity" are used in reference to nucleic acids (*i.e.*, a sequence of nucleotides) related by the well-known base-pairing rules that A pairs with T and C pairs with G. For example, the sequence 5'-A-G-T-3', is complementary to the sequence 3'-T-C-A-5'. Complementarity can be "partial," in which only some of the nucleic acid bases are matched according to the base pairing rules. On the other hand, there may be "complete" or "total" complementarity between the nucleic acid strands when all of the bases are matched according to base pairing rules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands as known well in the art. This is of particular importance in detection methods that depend upon binding between nucleic acids, such as those of the invention. The term "substantially complementary" refers to any probe that can hybridize to either or both strands of the target nucleic acid sequence under conditions of low stringency as described below or, preferably, in polymerase reaction buffer (Promega, M195A) heated to 95°C and then cooled to room temperature. As used herein, when the nucleic acid probe is referred to as partially or totally complementary to the target nucleic acid, that refers to the 3'-terminal region of the probe (*i.e.* within about 10 nucleotides of the 3'-terminal nucleotide position).

As used herein, the term "hybridization" is used in reference to the pairing of complementary

nucleic acid strands. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between nucleic acid strands) is impacted by many factors well known in the art including the degree of complementarity between the nucleic acids, stringency of the conditions involved affected by such conditions as the concentration of salts, the  $T_m$  (melting temperature) of the formed hybrid, the presence of other components (*e.g.*, the presence or absence of polyethylene glycol), the molarity of the hybridizing strands and the G:C content of the nucleic acid strands.

As used herein, the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds, under which nucleic acid hybridizations are conducted. With "high stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of complementary base sequences. Thus, conditions of "weak" or "low" stringency are often required when it is desired that nucleic acids which are not completely complementary to one another be hybridized or annealed together. The art knows well that numerous equivalent conditions can be employed to comprise low stringency conditions.

As used herein, the term " $T_m$ " is used in reference to the "melting temperature". The melting temperature is the temperature at which 50% of a population of double-stranded nucleic acid molecules becomes dissociated into single strands. The

equation for calculating the  $T_m$  of nucleic acids is well-known in the art. The  $T_m$  of a hybrid nucleic acid is often estimated using a formula adopted from hybridization assays in 1 M salt, and commonly used for calculating  $T_m$  for PCR primers:  $T_m = [(number\ of\ A + T) \times 2^\circ C + (number\ of\ G + C) \times 4^\circ C]$ . C.R. Newton et al. PCR, 2<sup>nd</sup> Ed., Springer-Verlag (New York: 1997), p. 24. This formula was found to be inaccurate for primers longer than 20 nucleotides.

*Id.* Other more sophisticated computations exist in the art which take structural as well as sequence characteristics into account for the calculation of  $T_m$ . A calculated  $T_m$  is merely an estimate; the optimum temperature is commonly determined empirically.

The term "homology," as used herein, refers to a degree of complementarity. There can be partial homology or complete homology (*i.e.*, identity). A partially complementary sequence that at least partially inhibits a completely complementary sequence from hybridizing to a target nucleic acid is referred to using the functional term "substantially homologous."

When used in reference to a double-stranded nucleic acid sequence such as a cDNA or genomic clone, the term "substantially homologous," as used herein, refers to a probe that can hybridize to a strand of the double-stranded nucleic acid sequence under conditions of low stringency.



When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous", as used herein, refers to a probe that can hybridize to (*i.e.*, is the complement of) the single-stranded nucleic acid template sequence under conditions of low stringency.

The term "interrogation position", as used herein, refers to the location of a given base of interest within a nucleic acid probe. For example, in the analysis of SNPs, the "interrogation position" in the probe is in the position that would be complementary to the single nucleotide of the target that may be altered from wild type. The analytical output from a method of the invention provides information about a nucleic acid residue of the target nucleic acid that is complementary to an interrogation position of the probe. An interrogation position is within about ten bases of the actual 3'-terminal nucleotide of the nucleic acid probe, although not necessarily at the 3'-terminal nucleotide position. The interrogation position of the target nucleic acid sequence is opposite the interrogation position of the probe, when the target and probe nucleic acids are hybridized.

The term "identifier nucleotide", as used herein, refers to a nucleotide whose presence is to be detected in a process of the invention to identify whether a depolymerization reaction has occurred. The particular application of a method of the invention affects which residues are considered an identifier nucleotide. For a method using ATP

detection (e.g. luciferase/luciferin or NADH)  
wherein, during analysis, all nucleotides released in  
the depolymerization are "converted" to ATP with an  
enzyme such as NDPK, all nucleotides released are  
5 identifier nucleotides. Similarly, for a method  
using absorbance detection that does not distinguish  
between nucleotides, all released nucleotides are  
identifier nucleotides. For a mass spectrometric  
detection wherein all the released nucleotides are  
10 analyzed, all released nucleotides can be identifier  
nucleotides; alternatively a particular nucleotide  
(e.g. a nucleotide analog having a distinctive mass)  
can be detected. For fluorescence detection, a  
fluorescently-labeled nucleotide is an identifier  
15 nucleotide. The nucleotide can  
be labeled, or the fluorescence level modified,  
prior to or after release from the nucleic acid. For  
radiographic detection, a radioactively-labeled  
nucleotide is an identifier nucleotide. In some  
20 cases, the release of identifier nucleotide is  
deduced by analyzing the remainder of the probe after  
a depolymerization step of the invention. Such  
analysis is generally by a determination of the size  
or mass of the remaining probe and can be by any of  
25 the described analytical methods (e.g. a fluorescent  
tag on the 5'-terminus of the probe to monitor its  
molecular weight following capillary  
electrophoresis).

The term "sample" is used in its broadest  
30 sense. A sample suspected of containing a nucleic  
acid can comprise a cell, chromosomes isolated from a

cell (e.g., a spread of metaphase chromosomes),  
genomic DNA, RNA, cDNA and the like.

The term "detection", as used herein,  
refers to quantitatively or qualitatively identifying  
5 a nucleotide or nucleic acid within a sample.

The term "depolymerization", as used  
herein, refers to the removal of a nucleotide from  
the 3' end of a nucleic acid.

The term "allele", as used herein, refers  
10 to an alternative form of a gene and the term  
"locus", as used herein, refers to a particular place  
on a nucleic acid molecule.

#### Detailed Description of the Invention

15 A contemplated method is utilized to assay  
for the presence or absence of nucleic acid that is  
exogenous to the source of the sample. For example,  
a contemplated method can be used to assay for the  
presence of viruses such as hepatitis C virus (HCV),  
20 cytomegalovirus (CMV), human immunodeficiency virus  
(HIV), as well as to determine the viral load in an  
organism with a disease, such as a human or a plant.  
A contemplated method can also be used to identify  
the presence of an exogenous nucleic acid sequence in  
25 a plant such as maize, soy or rice. A contemplated  
method can also be used to assay for the presence of  
microorganisms such as *Listeria monocytogenes*,  
*Campylobacter spp.*, *Salmonella spp.*, *Shigella spp.* or  
*Escherichia coli* (including *E. coli* E0157) in  
30 foodstuffs such as meats, dairy products, and fruit  
juices.

The determination of an appropriate exogenous nucleic acid target sequence useful for designing nucleic acid probes for use in a method of the invention is within the skill of the art.

5 Databases of genetic sequences, such as Genbank, can be used to ascertain the uniqueness of the selected nucleic acid target. Commercially available software for designing PCR primers can be used to assist in the design of probes for use in the invention.

10 A method of this invention is used to determine the presence or absence of at least one predetermined (known) exogenous nucleic acid target sequence in a nucleic acid sample. A nucleic acid target is "predetermined" in that its sequence must  
15 be known to design a probe that hybridizes with that target. However, it should be noted that a nucleic acid target sequence, as used with respect to a process of this invention, may merely act as a reporter to signal the presence of a different  
20 nucleic acid whose presence is desired to be determined. That other nucleic acid of interest does not have to have a predetermined sequence. Furthermore, a process of the invention is useful in determining the identity of base within a target  
25 where only enough of the sequence is known to design a probe that hybridizes to that exogenous target with partial complementarity at the 3'-terminal region of the probe.

30 Such a method utilizes an enzyme that can depolymerize the 3'-terminus of an oligonucleotide probe hybridized to the nucleic acid target sequence

to release one or more identifier nucleotides, under depolymerizing conditions, whose presence or absence can then be determined as an analytical output that indicates the presence or absence of the target sequence.

A nucleic acid target sequence is predetermined (or known) in that a nucleic acid probe is provided to be partially or totally complementary to that nucleic acid target sequence. A nucleic acid target sequence is a portion of nucleic acid sample with which the probe hybridizes if that target sequence is present in the sample.

A first step of the method is admixing a sample to be assayed with one or more nucleic acid probes. The admixing of the first step is typically carried out under low stringency hybridizing conditions to form a hybridization composition. In such a hybridization composition, the 3'-terminal region of the nucleic acid probe(s) (i) hybridizes with partial or total complementarity to an exogenous nucleic acid target sequence that may be present in the sample; and (ii) includes an identifier nucleotide in the 3'-terminal region.

Preferably, the nucleic acid probe is designed to not hybridize with itself to form a hairpin structure in such a way as to interfere with hybridization of the 3'-terminal region of the probe to the target nucleic acid. Parameters guiding probe design are well known in the art.

The hybridization composition is maintained under hybridizing conditions for a time period

sufficient to form a treated sample that may contain at least one predetermined nucleic acid target hybridized with a nucleic acid probe.

5 In the event that the sample to be assayed does not contain a target sequence to which the probe hybridizes, no hybridization takes place. When a method of the present invention is used to determine whether a particular target sequence is present or absent in a sample to be assayed, the resulting  
10 treated sample may not contain a substrate for the enzymes of the present invention. As a result, a 3' terminal region identifier nucleotide is not released and the analytical output is at or near background levels.

15 The treated sample is admixed with a depolymerizing amount of an enzyme whose activity is to release one or more identifier nucleotides from the 3'-terminal region of the probe that is hybridized to the nucleic acid target to form a  
20 depolymerization reaction mixture. The choice of enzyme used in the process determines if a match or mismatch at the 3'-terminal nucleotide results in release of that 3'-terminal nucleotide. Further information regarding specific enzyme reaction  
25 conditions is discussed in detail hereinafter.

The depolymerization reaction mixture is maintained under depolymerizing conditions for a time period sufficient to permit the enzyme to  
30 depolymerize hybridized nucleic acid and release identifier nucleotides therefrom to form a treated reaction mixture.

The presence or absence of released identifier nucleotides is then determined to obtain an analytical output. The analytical output indicates the presence or absence of at least the one nucleic acid target sequence.

Hybridization conditions can be empirically ascertained for a control sample for various time periods, pH values, temperatures, nucleic acid probe/target combinations and the like. Exemplary maintenance times and conditions are provided in the specific examples hereinafter and typically reflect low stringency hybridization conditions. In practice, once a suitable set of hybridization conditions and maintenance time periods are known for a given set of probes, an assay using those conditions provides the correct result if the nucleic acid target sequence is present. Typical maintenance times are about 5 to about 60 minutes.

The conditions and considerations with respect to hybridization of PCR primers to template nucleic acid in PCR are applicable to the hybridization of nucleic acid probes to target sequences in a process of the invention. Such hybridization conditions are well known in the art, and are a matter of routine experimentation depending on factors including the sequence of the nucleic acid probe and the target nucleic acid [sequence identity (homology), length and G+C content] molar amounts of nucleic acid present, buffer, salt content and duplex  $T_m$  among other variables.

Processes of the invention are sensitive and hybridization conditions of low stringency (e.g. temperature of 0-4°C) are sufficient, but moderate stringency conditions (e.g. temperatures of 40-60°C) also permit hybridization and provide acceptable results. This is true for all processes of the invention.

In one contemplated embodiment of the invention, the enzyme whose activity is to depolymerize hybridized nucleic acid to release nucleotides from the probe 3'-terminal end is a template-dependent polymerase. In such an embodiment, the reverse of a polymerase reaction is used to depolymerize a nucleic acid probe, and the identifier nucleotide is released when the 3'-terminal nucleotide of the nucleic acid probe hybridizes with total complementarity to its nucleic acid target sequence. A signal confirms the presence of a nucleic acid target sequence that has the sequence sufficiently complementary to the nucleic acid probe to be detected by the process of the invention.

In an embodiment that uses a 3'→5' exonuclease activity of a polymerase, such as Klenow or T4 DNA polymerase (but not limited to those two enzymes), to depolymerize a nucleic acid probe, an identifier nucleotide is released when the 3'-terminal residue of the nucleic acid probe is mismatched and therefore there is only partial complementarity of the 3'-terminus of the nucleic acid probe to its nucleic acid target sequence. In



this embodiment, to minimize background, the hybrid is typically purified from the unannealed nucleic acid prior to the depolymerization enzyme reaction, which may release an identifier nucleotide. A signal  
5 confirms the presence of a nucleic acid target sequence that is not totally complementary to the nucleic acid probe.

In an embodiment that uses a 3' → 5' exonuclease activity of Exonuclease III to  
10 depolymerize a nucleic acid probe, an identifier nucleotide is released when the 3'-terminal residue of the nucleic acid probe is matched to the target nucleic acid. A signal confirms the presence of a nucleic acid target that is complementary at the  
15 released identifier nucleotide.

It is thus seen that hybridization and depolymerization can lead to the release of an identifier nucleotide or to little or no release of such a nucleotide, depending upon whether the  
20 probe:target hybrid is matched or mismatched at the 3'-terminal region. This is also dependent on the type of enzyme used and the type of end, matched or mismatched, that the enzyme requires for depolymerization activity.

25 The magnitude of a contemplated analytical output under defined conditions is dependent upon the amount of released identifier nucleotides. Where an identifier nucleotide is released, an analytical output can be provided that has a value greater than  
30 background. Where an identifier nucleotide is not released either because the target sequence was not

present in the original sample or because the probe and depolymerizing enzyme chosen do not provide release of a 3'-terminal nucleotide when the target is present, or if the match/mismatch state of the 3'-  
5 terminal nucleotide did not match that required for the enzyme used to release a 3'-terminal nucleotide, the analytical output is substantially at a background level.

Contemplated methods and kits of the  
10 invention are useful for many applications as discussed above. For example, it is desirable to detect exogenous nucleic acid sequences when one would like to detect the presence or amount of viral contamination (or viral load) in a sample, often a  
15 biological or medical sample, but also in a food sample. The art provides many sequences that are useful for a wide variety of viral targets, and as more sequences are discovered, they are likewise useful in a process of the present invention. Thus,  
20 also contemplated are plant viruses, such as the tobacco mosaic virus. Exemplary viral probes follow.

Cytomegalovirus sequence probes.

SEQ ID NO:82 5'CGTTGTGCGGGTTCACGTCGATGAGCACGT  
25 TCATGGGTGTAATATCAAAGTGGCATAACGAGCT 3'

SEQ ID NO:35 5'CACTTTGATATTACACCCATG 3'

JH67 5' TCACACAGGAAACAGCTATGACCATG 3' SEQ ID NO:41

Hepatitis C virus probe.

HCV1:5' CTGCTAGCCGAGTAGTGTGGGTCGCGAAAGGCCTTGTGG 3'

5 SEQ ID NO:43

Human Immunodeficiency virus probes.

11808 5' CCATTTAGTACTGTCT 3' SEQ ID NO:52

11810 5' CTAGTTTTCTCCATTT 3' SEQ ID NO:54

10 11812 5' TTCTCTGAAATCTACT 3' SEQ ID NO:56

11814 5' AAAAAAGACAGTACTAAATGGAGAAAAGTAGTA  
GATTCAGAGAACTTAA 3' SEQ ID NO:58

15 Below are provided sequences of probes to  
various mutated viruses. Such probes are useful for  
distinguishing the presence of a particular mutated  
pathogen among other pathogens present.

Probes to a mutated cytomegalovirus sequences.

20 CV2 5' CACTTTGATATTACACCCGTG 3' SEQ ID NO:36

5' CGTGTATGCCACTTTGATATTACACCCGTGAACGTGCTCATCGACGTGAAC  
CCGCACAACGAGCT 3' SEQ ID NO:83

25 5' CGTTGTGCGGGTTCACGTCGATGAGCACGTTACGGGTGTAATATCAAAGT  
GGCATAACGAGCT3' SEQ ID NO:84

gancyclovir-resistant cytomegalovirus probes:

CV11 5' CGCTTCTACCACGAATGCTCGCAGACCATGCTGCACGAAT

30 ACGTCAGAAAGAACGTGGAGCGTCTGTTGGAGCT 3' SEQ ID NO:1

CV12 5' CCAACAGACGCTCCACGTTCTTTCTGACGTATTCGTGCAGC  
ATGGTCTGCGAGCATTTCGTGGTAGAAGCGAGCT 3' SEQ ID NO:2

CV13 5' CGCTTCTACCACGAATGCTCGCAGATCATGCTGCACGAAT  
5 ACGTCAGAAAGAACGTGGAGCGTCTGTTGGAGCT 3' SEQ ID NO:3

CV14 5' CCAACAGACGCTCCACGTTCTTTCTGACGTATTCGTGC  
AGCATGATCTGCGAGCATTTCGTGGTAGAAGCGAGCT 3' SEQ ID NO:4

10 Probes to drug-resistant HIV.

11815 5' AAAAAAACAGTACTAAATGGAGAAACTAGTAGA  
TTTCAGAGAACTTAA 3' SEQ ID NO:59

11816 AAAAAAGACAGTACTAGATGGAGAAACTAGTAGATTTTCAG  
15 AGAACTTAA 3' SEQ ID NO:60

11817 5' AAAAAAGACAGTACTAAATGGAGAAACTAA  
TAGATTTTCAGAGAACTTAA 3' SEQ ID NO:61

20 11813 5' TTCTCTGAAATCTATT 3' SEQ ID NO:57

11811 5' CTAGTTTCTCCATCT 3' SEQ ID NO:55

11809 5' CCATTTAGTACTGTTT 3' SEQ ID NO:53

25

The presence of other exogenous nucleic acids, such as those stemming from contaminating bacteria are useful in a process of the invention. Examples of bacterial probes follow.

30

Probes for *Listeria iap*.

LM1 5' GAAGTAAAACAAACTACACAAGCAACTACACCTGCGCCTAAAG  
TAGCAGAAACGAAAGAAACTCCAGTAG 3' SEQ ID NO:9

LM2 5' CTACTGGAGTTTCTTTCGTTTCTGCTACTTTAGGCGCAGGT  
5 GTAGTTGCTTGTGTAGTTTGTGTTTACTTC 3' SEQ ID NO:10

LM3 5' GCAACTACACCTGCGCCTAAAGTAGCAGAA 3' SEQ ID NO:11

LM4 5' TTCTGCTACTTTAGGCGCAGGTGTAGTTCG 3' SEQ ID NO:12

10 Probes for *Listeria hyl*.

LM5 5' CATCGACGGCAACCTCGGAGACTTACGAGATATTTTGAAAAAA  
GGCGCTACTTTTAATCGAGAAACACCA 3' SEQ ID NO:13

LM6 5' TGGTGTTTCTCGATTAAAGTAGCGCCTTTTTTCAAAATATCT  
15 CGTAAGTCTCCGAGGTTGCCGTCGATG 3' SEQ ID NO:14

LM7 5' CTCGGAGACTTACGAGATATTTTGAAAAAA 3' SEQ ID NO:15

LM8 5' TTTTTTCAAAATATCTCGTAAGTCTCCGAG 3' SEQ ID NO:16

20

Probes for *Salmonella*.

ST3 5' TGTGTAATGAAAGAAATCACCGTCACTGAA 3' SEQ ID NO:19

ST4 5' TTCAGTGACGGTGATTTCTTTTATTACACA 3' SEQ ID NO:20

25 Probes to *Campylobacter jejuni*.

11453 5' CTTGAAGCATAGTTCTTGTTTTTAACTTTGTCCATCTT  
GAGCCGCTTGAGTTGCCTTAGTTTTTAATAGT 3'

SEQ ID NO:31

11454 5' ACTATTAAACTAAGGCAACTCAAGCGGCTCAAGATGG

30 ACAAAGTTTAAAAACAAGAACTATGCTTCAAG 3'

SEQ ID NO:33

11451 5'AGTTCTTGTTTTTAACTTTGTCCATCTTG 3'

SEQ ID NO:32

11450 5'CAAGATGGACAAAGTTTAAAAACAAGAACT 3'

SEQ ID NO:34

5

It is often desirable to detect the presence of exogenous genes, typically incorporated merely as markers for inserted genes. Classic molecular biology techniques include the  
10 incorporation of antibiotic resistance to select for clones having the desired inserted exogenous nucleic acid sequence. The antibiotic resistance gene is also an exogenous sequence. The sequences of  
exogenous "marker" genes are well-known in the art  
15 and are easily available to a worker of ordinary skill. Exemplary fragments of such sequences useful as probes in methods and kits of the present invention follow.

20 Probes for genes conferring kanamycin resistance to bacteria.

5'GCAACGCTACCTTTGCCATGTTTC 3' SEQ ID NO:21

5'GCAACGCTACCTTTGCCATGTTTG 3' SEQ ID NO:22

5'GCAACGCTACCTTTGCCATGTTTA 3' SEQ ID NO:23

25 5'GCAACGCTACCTTTGCCATGTTTT 3' SEQ ID NO:24

5'GCAACGCTACCTTTGCCATGTTTC 3' SEQ ID NO:85

Probes to the  $\beta$ -galactosidase gene, commonly used as a biological marker for exogenous genes in the field  
30 of biochemistry.

5'CAGTCACGACGTTGTAAAACGACGGCCAGT3' SEQ ID NO:29

5'ACTGGCCGTCGTTTTACAACGTCGTGACTG3' SEQ ID NO:30

Other exogenous sequences, though not necessarily used as markers, are useful in methods and kits of the present invention. This is discussed further in the examples below, particularly with respect to genetically modified organisms. Some exemplary common exogenous sequences that are often introduced along during genetic engineering of an organism follow.

Probes for the plant 35S promoter commonly used in biotechnology when inserting exogenous plant genes.

11211	5' GCAAGTGGATTGATG 3'	SEQ ID NO:48
11210	5' CCAACCACGTCTTCAAA 3'	SEQ ID NO:49

Probes for the plant NOS terminators commonly used in biotechnology when inserting exogenous genes.

11212	5' TTTATGAGATGGGTTT 3'	SEQ ID NO:50
11213	5' ATGATTAGAGTCCCG 3'	SEQ ID NO:51

In one embodiment of the invention, viral load, the amount of virus present, is determined from the magnitude of the analytical output from a predetermined amount of biological sample such as a animal fluid or tissue. Processes of the invention are quantitative and very sensitive. The sensitivity is enhanced further through use of a process of the invention including a step to enrich the sample in the predetermined exogenous nucleic acid target sequence, by conversion of a signal from the

predetermined exogenous nucleic acid target sequence  
to an amplified predetermined reporter sequence or by  
amplification of the signal from the released  
identifier nucleotide. In the viral load example  
5 below, the target sequence is enriched in the sample  
through RT-PCR.

In one aspect of the method, the  
predetermined nucleic acid target sequence is present  
in the sample for the purpose of gene therapy. An  
10 exemplary gene therapy embodiment would be in the  
provision of an exogenous gene to an animal,  
preferably a person or commonly raised animal such as  
a cow, pig, sheep, dog or chicken, to make up for a  
missing gene, such as is the case with phenyl  
15 ketonuria or persons lacking adenine deaminase.

A worker of ordinary skill in the art  
recognizes that processes and kits of the invention  
are useful with any predetermined sequence that is  
specifically sought for assay. Such a worker need  
20 only construct a nucleic acid probe that is  
complementary to the predetermined sequence. Thus  
the present invention is useful for determining the  
success of genetic engineering into a plant,  
typically a crop, by searching for the introduced  
25 gene. Similarly, the success of plant breeding is  
monitored using a process of the invention when the  
gene sought to be introduced into the cross-bred  
generation is a nucleic acid target.

A worker of ordinary skill further  
30 recognizes that it is possible to construct any  
desired probes for the specific methods using the



invention disclosed in the parent application, U.S. Serial No. 09/358,972, filed on July 21, 1999. (This application is a continuation-in-part of U.S. patent application Serial No. 09/358,972, the disclosures of which are herein by reference, and published on the internet at <http://www.promega.com/pt/pend/09358972.pdf>)

In one embodiment, the identity of a specific base located at an interrogation position within a nucleic acid target sequence is determined. This is useful for determination of the presence or absence of certain mutations within a gene. Examples of such mutated genes were listed above, such as the resistant HIV, or the gancyclovir resistant CMV.

#### **Depolymerization**

Depolymerization reactions and enzymes useful in such reactions are discussed below. Nucleic acid polymerases generally catalyze the elongation of nucleic acid chains. The reaction is driven by the cleavage of a pyrophosphate released as each nucleotide is added. Each nucleoside-5'-triphosphate has three phosphate groups linked to carbon five of the ribose or deoxyribose sugar. The addition of a nucleotide to a growing nucleic acid results in formation of an internucleoside phosphodiester bond. This bond is characterized in having a 3' linkage to carbon 3 of ribose or deoxyribose and a 5' linkage to carbon 5 of ribose or deoxyribose. Each nucleotide is added through formation of a new 3' to 5' linkage, so the nucleic acid strand grows in a 5' to 3' direction.

Depolymerization in its strictest sense means the reverse of polymerization so that in the present context, an internucleotide phosphodiester bond is broken between the two 3'-terminal bases in

the presence of pyrophosphate and a polymerase enzyme to form a nucleic acid that is one nucleotide shorter and a nucleoside triphosphate. A somewhat more encompassing definition is contemplated here. In accordance with that definition, the 3'-terminal nucleotide is removed from a nucleic acid in a reaction catalyzed by an enzyme, but the nucleotide formed can be a monophosphate and pyrophosphate is not always required.

10           The former reactions (i.e. reverse of polymerization) are referred to herein as pyrophosphorolysis reactions whereas the latter, more encompassing definition, reactions are referred to as exonuclease reactions. It is to be understood that the depolymerization reaction of interest in the invention is that depolymerization occurring in the 3'-terminal region of the nucleic acid probe. This depolymerization reaction releases identifier nucleotides under appropriate depolymerizing conditions, as discussed herein.



20           Depolymerization reactions and enzymes useful in such reactions are discussed in parental U.S. Patent Application Serial No. 09/358,972, filed on July 21, 1999, which disclosure is incorporated herein by reference.

#### **A. Pyrophosphorolysis**



30           In some embodiments of the present invention, a method comprises depolymerizing the nucleic acid (NA) at a 3'-terminal nucleotide by enzymatically cleaving the terminal internucleoside

phosphodiester bond in the presence of pyrophosphate, or an analogue thereof, to form an XTP (e.g. NTPs or dNTPs) as illustrated by the following reaction on double-stranded DNA having a 5' overhang:

5



10



15

Template-dependent nucleic acid polymerases capable of pyrophosphorolysis include, but are not limited to, DNA polymerase  $\alpha$ , DNA polymerase  $\beta$ , T4 DNA polymerase, *Taq* polymerase, *Tne* polymerase, *Tne* triple mutant polymerase, *Tth* polymerase, *Tvu* polymerase, *Ath* polymerase, *Bst* polymerase, *E. coli* DNA polymerase I, Klenow fragment, Klenow exo minus (exo-), AMV reverse transcriptase, RNA polymerase and MMLV reverse transcriptase, and poly(A) polymerase.

20

25

Most preferably, Klenow exo minus (Klenow exo-) or *Tne* triple mutant polymerase is utilized for DNA pyrophosphorolysis reactions because of their efficient utilization of 5' overhanging DNA ends.

30

In a preferred embodiment in the case of the reverse of polymerase activity (pyrophosphorolysis), a preferred substrate is a DNA probe hybridized to an exogenous nucleic acid target sequence with total complementarity at its 3'-terminus, including an identifier residue in the 3'-terminal region. In an example of this preferred

embodiment, when the nucleic acid probe is hybridized to an exogenous nucleic acid target sequence such that there is one base mismatch at the 3'-terminal nucleotide of the nucleic acid probe, the nucleic acid probe is inefficiently depolymerized through the reverse polymerization reaction. Thus, such a substrate is not an ideal substrate for depolymerization.

The non-ideality of the substrate for depolymerization via a reverse of the polymerization reaction is recognized with a single base mismatch as far in as about 10 residues from the 3'-terminus of the nucleic acid probe. With a single base mismatch 12 residues from the 3'-terminus of the probe, the depolymerization reaction can occur to approximately the same extent as when there is no mismatch and the nucleic acid probe is totally complementary to the nucleic acid target sequence.

It is thus contemplated that the reactivity of the depolymerization reaction is a continuum that is related to the efficiency of the substrate. A partially complementary hybrid is a less efficient depolymerization substrate than a totally complementary hybrid for the reverse of a polymerization reaction. It is contemplated that this differential reactivity be used to enhance the discrimination between matches and mismatches at certain positions (e.g. an interrogation position). When a substrate hybrid is totally complementary, it will give a fairly high analytical output. A mismatch can be intentionally introduced to

destabilize the substrate hybrid. Such a destabilization can increase the difference in analytical output between bases substituted at an interrogation position that is different from the destabilizing base position.

Several chemical compounds are known in the art to be substitutable for pyrophosphate in pyrophosphorolysis reactions. Rozovskaya, et al., *Biochem. J.*, **224**:645-650 (1984).

Preferred reaction mixtures and times (depolymerization conditions) for depolymerization by pyrophosphorolysis, including suitable buffers for each nucleic acid polymerase analyzed, are described in greater detail in the Examples. Typically, under these conditions, sufficient NTP or dNTP is released to accurately detect or assay extremely low amounts of nucleic acids (e.g., about 5-1000 picograms). ATP can be produced by conversion from XTP by an enzyme such as NDPK (in the presence of ADP) prior to analysis or the ATP can be further amplified prior to analysis.

The high efficiency of the pyrophosphorolysis reaction was unexpected, and appears to be associated with extremely low levels of DNA substrate, in contrast to previous DNA pyrophosphorolysis studies conducted using much greater amounts of DNA.

The pyrophosphorolysis activity of different nucleic acid polymerases also varies. For example, T4 polymerase and *The* DNA polymerase possess very high pyrophosphorolysis activity as measured by

a luciferase assay for ATP produced by  
pyrophosphorolysis. Pyrophosphorolysis using T4  
polymerase resulted in about a 10 fold increase in  
light production as compared to MMLV-RT and a 4-fold  
5 increase in light production as compared to *Taq*  
polymerase.

The type of DNA end resulting from  
restriction enzyme digestion also affects the  
pyrophosphorolysis activity of different nucleic acid  
10 polymerases. For example, Klenow exo-, MMLV-RT and  
*Taq* polymerase catalyze pyrophosphorolysis of DNA  
fragments with 5'-overhangs and with blunt-ends, but  
have little or no pyrophosphorolysis activity with  
3'-overhangs. In contrast, T4 DNA polymerase  
15 catalyzes both 3'- and 5'-end overhang and blunt-end  
mediated pyrophosphorolysis. Thus, T4 DNA polymerase  
is a preferred enzyme for pyrophosphorolysis of a  
hybrid with a 3'-overhang. When other nucleic acid  
polymerases are utilized for pyrophosphorolysis of  
20 restriction enzyme treated DNA, it is contemplated  
that care is taken to match the end specificity of  
the polymerase with the type of end created by the  
restriction endonuclease. Such care is well within  
the skill of those in the art.

25 Further, it is contemplated that the type  
of polymerase used in the pyrophosphorolysis reaction  
is matched to the correct nucleic acid substrate in  
order to produce the best results. In general, DNA  
polymerases and reverse transcriptases are preferred  
30 for depolymerizing DNA, whereas RNA polymerases are  
preferred for depolymerizing RNA. Reverse

transcriptases or DNA polymerases with reverse transcriptase activity are preferred for depolymerizing RNA-DNA hybrids.

5 In the grandparent application, it was surprisingly demonstrated that poly(A) polymerase can catalyze pyrophosphorolysis, even though no such reaction had been previously reported. Indeed, poly(A) polymerase has been widely reported to not catalyze pyrophosphorolysis. (See e.g., Sippel, *Eur. J. Biochem.*, 37:31-40 (1973) and Sano and Feix, *Eur. J. Biochem.*, 71:577-83 (1976)). However there are many differences between the conditions used in the grandparent application disclosure and those reported in the references. In these preferred embodiments of 10 the invention disclosed in the grandparent application, the manganese chloride present in the previously reported buffers is omitted, the concentration of sodium chloride is decreased, and the pH value is lowered from about 8.0 to about 7.5. 15 Furthermore, the poly(A) polymerase pyrophosphorolysis reaction buffer contains about 50 mM Tris-Cl pH 7.5, 10 mM MgCl<sub>2</sub>, 50 mM NaCl, and 2 mM NaPP<sub>i</sub> (sodium pyrophosphate). 20

It is important to note that the 25 depolymerization reaction is the reverse of the polymerization reaction. Therefore, as increasing amounts of free nucleoside triphosphates are produced by depolymerization, a state of equilibrium can theoretically be attained in which polymerization and 30 depolymerization reactions are balanced. Alternatively, where small amounts of nucleic acid

are detected, the reaction can go essentially to completion without reaching equilibrium, (i.e., the nucleic acid target is depolymerized into its constituent subunit nucleotides by greater than 50%).

5 This factor is important in quantitative assays because the total amount of nucleotides released is proportional to the amount of signal generated in the detection assay.

When used for qualitative detection of  
10 nucleic acid, as long as a threshold level of nucleotides is produced, it is not necessary that the reaction reach equilibrium or go essentially to completion. In preferred embodiments, the mixture of nucleoside triphosphate molecules produced by  
15 depolymerization is preferably converted to ATP as described below. For either quantitative or qualitative detection, a detectable threshold ATP concentration of approximately  $1 \times 10^{-12}$  molar in 100  $\mu$ l of sample is preferably provided for detection of  
20 light in a typical luciferase assay.

In some preferred embodiments, oligonucleotide probes are typically utilized at about 100 ng to about 1  $\mu$ g per 20  $\mu$ L depolymerization reaction. That amount provides a probe to target  
25 weight ratio of about 200:1 to about 1,000:1.

In a preferred embodiment of the present invention, nucleic acid polymerase and pyrophosphate ( $PP_i$ ) or an analogue thereof, are added to a hybridized sample containing from less than about 100  
30  $\mu$ g of target nucleic acid, to less than about 10 pg of nucleic acid. Typical target nucleic acids are



present at about 1 to about 5 ng in the sample to be assayed, with a target nucleic acid length of about 30 to about 1000 bp being preferred.

When using enzymes that utilize 5' overhang substrates, it is preferred that the 3' end of the target nucleic acid extends beyond the 5' end of the nucleic acid probe. In this way, the only 5' overhang substrate is that where the 5' end of the target nucleic acid overhangs the 3' terminal region of the nucleic acid probe. An alternative method of limiting depolymerization to the nucleic acid probe is chemical modification of the ends of other nucleic acids in the sample, such as, for example, making a phosphorothioate linkage at the 3'-terminus of the target nucleic acid.

A depolymerizing enzyme is preferably present in an amount sufficient to depolymerize a hybridized target:probe. That amount can vary with the enzyme used, the depolymerization temperature, the buffer, and the like, as are well-known in the art. For a typical reaction carried out in a 20  $\mu$ L volume, about 0.25 to about 1 unit (U) of an enzyme such as Klenow exo- is used. About 1 to about 5 U of the thermostable enzymes are used for depolymerization at elevated temperatures.

Luciferase, which is part of the preferred ATP detection system, is inhibited by  $PP_i$ . In preferred embodiments, care is taken to avoid transferring a highly inhibiting amount of  $PP_i$  to the ATP detection reaction. Preferably, the amount of  $PP_i$  carried over to the ATP detection reaction results in

a concentration of  $PP_i$  in the luciferase detection reaction of less than about  $100\ \mu M$ , although less than about  $10\ \mu M$  is desirable. Therefore, the amount of  $PP_i$  utilized in the pyrophosphorolysis reaction is determined by the size of the aliquot that is taken for use in the luciferase detection system. It is contemplated that the aliquot size can vary depending upon the test system used, but the amount of  $PP_i$  transferred or carried over to the luciferase detection reaction corresponds to the  $PP_i$  concentration parameters described above, so that the concentration of  $PP_i$  is at least below about  $100\ \mu M$ , and preferably below about  $10\ \mu M$ .

In one preferred embodiment of the invention, the enzyme whose activity is to depolymerize is a template-dependent polymerase. The depolymerization reaction is a reverse of the polymerization reaction. In a contemplated embodiment, the polymerization reaction is reversed in the presence of pyrophosphate in a reaction referred to as pyrophosphorolysis.

In some preferred embodiments, the reaction conditions are preferably adjusted to further favor depolymerization of a nucleic acid probe that is hybridized with its target nucleic acid sequence by providing a higher concentration of nucleic acid probe than its target nucleic acid sequence.

One strategy to favor the depolymerization of a probe:target hybrid is that the probe be in molar excess over the nucleic acid target in the

hybridization step after denaturing of duplex target nucleic acid.

Another strategy to favor the depolymerization of a probe:target hybrid is to isolate only the strand of nucleic acid target to which the probe is complementary. There are several techniques that can be used to achieve this end.


In one such technique, phosphorothioate linkages are utilized at the 5'-terminus of a target nucleic acid amplifying primer sequence, e.g., at the 1 to about 10 5'-most residues. Upon PCR amplification of the target, the phosphorothioate linkages of the primer become incorporated into the amplified target nucleic acid as part of one of a pair of complementary strands. Treatment of the double-stranded resulting molecule with T7 polymerase exonuclease 6 removes the non-phosphorothioate-containing strand.

In another technique, strand isolation can be accomplished by amplifying the target nucleic acid using PCR primers incorporated into the extended nucleic acid strand (with which a nucleic acid probe useful herein is designed to hybridize) that are not labeled, whereas primers for the complementary strand are labeled, such as with biotin. Then, the amplified nucleic acid is denatured and added to streptavidin linked to a solid support. A useful material is Streptavidin MagneSphere® paramagnetic particles (Promega, Z548A), where a magnet can be used to separate the desired target nucleic acid strand from its biotinylated complementary strand.

Further discussion pertaining to pyrophosphorolysis is found in the parent cases cited above and incorporated herein by reference.

5           **B. Exonuclease Digestion**

          In other embodiments of the present invention, a method comprises depolymerizing the nucleic acid at a 3'-terminal nucleotide by enzymatically cleaving the terminal internucleoside phosphodiester bond to form an XMP as illustrated by  
10           the following reaction on double-stranded DNA having a 5'-overhang:




          For example, such a hydrolysis reaction can be catalyzed by Klenow or Exonuclease III.

25           In some embodiments (e.g., quantitative assays for nucleic acids), the depolymerizing step is repeated essentially to completion or equilibrium to obtain at least two nucleotide molecules from a strand of minimally three nucleotides in order to  
30           increase detection sensitivity. In alternative embodiments, (e.g., qualitative detection of DNA), the depolymerizing step need not be repeated if there


are sufficient nucleic acid molecules present to generate a signal.

In another embodiment of the present invention, terminally mismatched hybridized nucleic acid probes are first depolymerized into NMP or dNMP by exonuclease digestion according to the following reaction:




wherein  $\text{NA}_n$  is a nucleic acid, XMP is either a dNMP or NMP, and  $n$  is the number of nucleotides in the nucleic acid.

This depolymerization reaction is shown more specifically below in the following reaction on double-stranded DNA having a 5'-overhang and mismatched bases at the 3'-terminus:



↓ enzyme




For example, such a depolymerization reaction can be catalyzed by bacteriophage T4 polymerase in the absence of NTPs. In preferred embodiments, the released nucleotides, XMPs, are produced by nuclease digestion.

Nuclease digestion can be accomplished by a variety of nucleases that release a nucleotide with a 5' phosphate, including S1 nuclease, nuclease Bal 31, mung bean nuclease, exonuclease III and ribonuclease H. Nuclease digestion conditions and buffers are known in the art. Nucleases and buffers for their use are available from commercial sources.

In the biosynthesis of purine and pyrimidine mononucleotides, phosphoribosyl-1-pyrophosphate (PRPP) is the obligatory ribose-5'-phosphate donor. PRPP itself is formed in a reaction catalyzed by PRPP synthetase through the transfer of pyrophosphate from ATP to ribose-5'-phosphate. This reaction is known to be reversible as described in Sabina *et al.*, *Science*, 223:1193-95 (1984).

In some embodiments of the present invention, the NMP or dNMP produced by nuclease digestion is preferably converted directly to NTP or dNTP by the enzyme PRPP synthetase in the following reaction:



wherein XMP is either AMP or dAMP, and XTP is either ATP or dATP. Preferably, this reaction produces a threshold ATP concentration of approximately  $1 \times 10^{-12}$  M in 100  $\mu$ l of sample.

In this reaction, the pyrophosphate group of PRPP is enzymatically transferred to XMP molecules, forming XTP molecules. Examples of

suitable reaction conditions and buffers are set forth elsewhere herein.

Utilization of the PRPP reaction in the nucleic acid detection system of the present invention has advantages over previously reported methods. For example, only one step is necessary to convert an AMP or dAMP to ATP or dATP, thereby simplifying the detection system. In addition, contamination of the detection reaction with exogenous ATP, ADP, or AMP is less likely using methods of the present invention, as compared to previously reported methods.

In an embodiment wherein the depolymerizing enzyme exhibits 3'→5' exonuclease activity, the substrate is a double-stranded or single-stranded nucleic acid having a 3'-hydroxyl terminus. Enzymes having 3'→5' exonuclease activity that are useful in a process of the invention include *E. coli* DNA polymerase I, Klenow fragment and bacteriophage T4 DNA polymerase. *E. coli* DNA polymerase I holoenzyme is not preferred in a process of the invention because it is preferable to avoid the 5'→3' exonuclease activity that degrades probe:target hybrids regardless of the degree of hybridization at the 3'-terminus. Bacteriophage λ exonuclease has only 5'→3' exonuclease activity, so it is not a contemplated enzyme. Similarly, *Taq* DNA polymerase has a very low level of 3'→5' exonuclease activity. Exonuclease III (Exo III) has 3' exonuclease activity on blunt-ended substrates or those having 5'-

overhangs or nicks with 3'-hydroxyl groups, and is thus useful in a process of the invention for depolymerizing hybrids with matched 3' terminal nucleotides. However, Exo III is not limited to  
5 hybrids having only partially complementary 3'-termini, it requires a double stranded end, i.e. a matched terminal nucleotide.

In an embodiment of the invention where the enzyme's activity is a 3'→5' exonuclease activity,  
10 the hybridized nucleic acid probe is depolymerized from its 3'-terminal nucleotide. In a preferred embodiment in the case of a 3'→5' exonuclease activity of a polymerase, the preferred substrate is a nucleic acid probe hybridized to an exogenous  
15 nucleic acid target sequence with partial complementarity at its 3'-terminal region, most preferably with a mismatch at its 3'-terminal residue that is an identifier nucleotide.

A contemplated method is particularly  
20 useful in a multiplex assay environment in which a plurality of probes is utilized to determine whether one or more of a plurality of predetermined exogenous nucleic acid sequences is present or absent in a sample. A particularly useful area for such  
25 multiplex assays is in screening assays where the usual analytical output indicates that the sought-after exogenous gene is absent.

In one illustrative embodiment, a nucleic acid sample is screened for the presence of a  
30 plurality of predetermined exogenous genes, e.g. viruses in a biological sample. In this embodiment,



the viruses usually are not present and the analytical output is, for example, at about background levels except where a virus with its exogenous nucleic acid is present.

5           In another embodiment, a plurality of samples is examined for the presence or absence of microbe-specific genes. Here, again, where a population of healthy individuals, animals, or presumably sterile food is sampled, the absence of  
10 the sought-after exogenous genes provides an analytical output that is about background levels, and only in the rare instance of microbial contamination does a greater than the background output appear.

15           In a multiplexed embodiment of the above process, the sample is admixed with a plurality of different exogenous nucleic acid probes, in some embodiments after amplification of the multiple nucleic acid targets as needed. In this embodiment  
20 of the invention, the analytical output for a certain result with one of the probes is distinguishable from the analytical output from the opposite result with all of the probes.

          In preferred embodiments, the ATP produced  
25 via NDPK conversion of released nucleotides in the presence of ADP is detected by a luciferase detection system or an NADH detection system. In still another embodiment of the present invention, the pyrophosphate transferring step and the phosphate  
30 transferring step are performed in a single pot reaction. In other preferred embodiments, if

increased sensitivity is required, the ATP molecules can be amplified.

#### **Analytical Output**

5           The analytical output is obtained by  
detection of the released identifier products, either  
the released nucleotides or the remainder of the  
probe. Exemplary detection systems include the light  
emitting luciferase detection system, the NADH light  
10   adsorption detection system (NADH detection system),  
fluorescence emissions and mass spectrometry. These  
detection systems are discussed hereinbelow.

          The fact that nucleotides were released (a  
qualitative determination), or even the number of  
15   nucleotides released (a quantitative determination)  
can be deduced through examination of the probe after  
depolymerization. The determination of the size of  
an oligonucleotide is well known in the art. For  
example gel separation and chromatographic  
20   separations are well known. Gel imaging techniques  
that take advantage of fluorescence and absorbance  
spectroscopy as well as radiographic methods. Mass  
spectrometry of oligonucleotides is also becoming  
more common.

25


#### **A. Luminescence spectroscopy**

          Luciferase detection systems are  
particularly useful for detecting ATP. In the  
presence of ATP and oxygen, luciferase catalyzes the  
30   oxidation of luciferin, producing light that can then  
be quantified using a luminometer. Additional

products of the reaction are AMP, pyrophosphate and oxyluciferin.

In particularly preferred embodiments, ATP detection buffer referred to as L/L reagent (Promega, FF2021) is utilized. Preferably, about 5 to 10 ng of luciferase are used in the reaction. Although it is not intended that the present invention be limited to a specific concentration of luciferase, greater amounts of luciferase have a tendency to increase non-specific background.

It is contemplated that in some embodiments, the dNTPs or NTPs produced by pyrophosphorolysis or nuclease digestion are converted to XTP, which can then be used directly as substrate for luciferase, permitting detection of the nucleic acid. However, the preferred substrate for luciferase is ATP, as demonstrated by Moyer and Henderson, *Anal. Biochem.*, 131:187-89 (1983). When DNA is the initial substrate, NDPK is conveniently utilized to catalyze the conversion of dNTPs to ATP by the following general reaction:



wherein dNTP is a mixture of deoxyribonucleoside triphosphates and dNDP is the corresponding deoxyribonucleoside diphosphate. In Reaction 3, the terminal 5'-triphosphate (P\*) of the dNTP is transferred to ADP to form ATP.

Enzymes catalyzing this reaction are generally known as nucleoside diphosphate kinases (NDPKs). NDPKs are ubiquitous, relatively

nonspecific enzymes. For a review of NDPK, see Parks and Agarwal, in *The Enzymes*, Volume 8, P. Boyer Ed. (1973).

5 The conversion of NTPs or dNTPs to ATP by NDPK is preferably accomplished by adding NDPK and a molar excess of ADP over the amounts of NTPs or dNTPs expected to be produced by pyrophosphorolysis or nuclease digestion, followed by pyrophosphorylation by PRPP synthetase. The utilization of ADP requires  
10 optimization of the amount of ADP added. Too much ADP results in high background levels.

NDPK (EC 2.7.4.6) preparations from several biological sources are commercially available from several suppliers. For example yeast NDPK is  
15 available from Sigma Chemical Co., St. Louis, MO, whereas bovine NDPK is available from ICN Biochemicals, Inc., Costa Mesa, CA. The particular NDPK selected for most uses described herein is typically a matter of choice.

20 The *Tne* triple mutant DNA polymerase is described in detail in WO 96/41014, whose disclosures are incorporated by reference, and its 610 residue amino acid sequence is provided as SEQ ID NO:35 of that document. That enzyme is referred to in WO  
25 96/41014 as *Tne* M284 (D323A,D389A).

Briefly, that enzyme is a triple mutant of the polymerase encoded by the thermophilic eubacterium *Thermotoga neapolitana* (ATCC 49049). The amino-terminal 283 residues of the native sequence  
30 are deleted and the aspartic acid residues at positions 323 and 389 of the native sequence are

replaced by alanine residues in this recombinant enzyme. This recombinant enzyme is thus a deletion and replacement mutant of the native enzyme.

Deletion of the amino-terminal sequence  
5 removes the 5' exonuclease activity of the native enzyme, whereas replacement of the two aspartic acid residues removes a magnesium binding site whose presence facilitates exonuclease activity, and this triple mutant also exhibited no 3' exonuclease  
10 activity relative to the recombinant native enzyme. This triple mutant enzyme exhibited a half-life at 97.5°C of 66 minutes as compared to the full length recombinant enzyme that exhibited a half-life of only 5 minutes at that temperature.

15 A reaction containing NDPK contains about 0.01 to 0.50  $\mu$ M ADP, preferably about 0.05  $\mu$ M ADP. Various useful buffers and other reaction components are set forth elsewhere. NDPK is itself present in an amount sufficient to catalyze the desired  
20 conversion of ADP to ATP. In a typical assay starting from a 20  $\mu$ L depolymerization reaction, about 0.1 U of NDPK are used.

Where larger volumes of reactants are used, with the target and probe concentrations being  
25 approximately proportionately larger, the amount of NDPK or the other enzymes discussed herein can be used in a similar larger proportion relative to the amount discussed for the 20  $\mu$ L reaction. Indeed, a 20  $\mu$ L reaction has been successfully scaled down

about two fold and scaled upwardly by a factor of about 20.

#### **B. Mass Spectrometric Analysis**

5           In one method of the invention, the presence of released nucleotides is analyzed via mass spectrometry. In an embodiment of a method using mass spectrometry, the treated reaction mixture is ionized in a manner such that all components of the  
10   treated reaction mixture in the molecular weight range of the released identifier nucleotides are measured. Very small differences in molecular weight can be detected using mass spectrographic methods (different isotopes of the same atom are detectable),  
15   so any variation from a natural nucleic acid, including a single atom substitution (e.g. a fluorine in place of a hydrogen atom or a replacement of a hydrogen by a deuterium atom) in the identifier nucleotide gives rise to a detectable difference.  
20   Nucleic acid analogs used in methods of the invention should not interfere with either the hybridization of the nucleic acid probe or depolymerization of the hybridized probe.

          Additionally, mass spectrometry can  
25   discriminate between individual nucleotides or nucleosides. For example, if the 3'-identifier nucleotide used in the instant invention was a G nucleotide, mass spectrometry can be used to detect the release of that G nucleotide in a method of the  
30   present invention. Similarly, mass spectrometry can detect the release of an A, T or C nucleotide, based

on the differences in atomic weight of these compounds. Thus, in a multiplexing embodiment of the present invention, mass spectrometry can be used to resolve the presence of one or more of these 3'-  
5 identifier nucleotides.

In a particularly useful aspect of this embodiment, a mass spectral technique referred to as DIOS (desorption/ionization on silicon) was recently reported by Wei et al., *Nature*, 399:243(1999) that  
10 can accurately perform one or multiple assays on picogram or attogram amounts using commercially available mass spectrographs adapted with a specialized porous silicon sample well. The older, well known, MALDI mass spectrographic assay  
15 techniques can also be utilized.

In an embodiment of a multiplex method using mass spectrometry, multiple different identifier nucleotides can be used in the various nucleic acid probes. Using such a technique the  
20 presence of the different identifier nucleotides is direct evidence of the presence of the nucleic acid target sequences.

### **C. Fluorescence Spectroscopic Analysis**

25 In some contemplated embodiments, the identifier nucleotide emits fluorescence. For example, in one embodiment when the nucleotide has a fluorescent label, the analytical output is obtained by fluorescence spectroscopy. In an alternative  
30 embodiment when the nucleotide has a fluorescent

label, the analytical output is obtained by mass spectrometry.

In a preferred embodiment of the invention, the fluorescent label is part of a fluorescent analog of a nucleotide. Fluorescent nucleotide analogs are widely known and commercially available from several sources. An exemplary source is NEN<sup>TM</sup> Life Science Products (Boston, Massachusetts), who offer dideoxy-, deoxy-, and ribonucleotide analogs labeled with fluorescein, coumarin, tetramethylrhodamine, naphthofluorescein, pyrene, Texas Red<sup>®</sup>, and Lissamine<sup>TM</sup>. Other suppliers include Amersham Pharmacia Biotech (Uppsala, Sweden; Piscataway, New Jersey) and MBI Fermentas, Inc. (Amherst, New York).

An advantage to using fluorescent labels and fluorescence spectroscopy analysis is that there are multiple different labels. Such different labels would be particularly useful in a multiplex embodiment of the invention. Different fluorescent labels would be used in different probes, so that the detection of a particular fluorescently-labeled nucleotide analog as a released identifier nucleotide could be used to deduce which nucleic acid targets are present.

For example, fluorescein has a 488 nm excitation and 520 nm emission wavelength, whereas rhodamine (in the form of tetramethyl rhodamine) has 550 nm excitation and 575 nm emission wavelength. A fluorescence detector provides an excitation source and an emission detector. The emission wavelengths



of 520 nm and 575 nm are easily distinguishable using fluorescence spectroscopy.

On a per molecule basis, fluorescence spectroscopy is about 10-fold more sensitive than absorbance spectroscopy. A very wide variety of fluorescence spectroscopy-based detectors are commercially available for reading fluorescence values of single tubes, flow cells and multi-well plates, among others. For example, Labsystems Multiskan models of microplate readers are widely available with a spectral range of 400 to 750 nm, and filters for 340, 405, 414, 450, 492, 540, 620, and 690 nm (e.g. Fisher Scientific, Pittsburgh, Pennsylvania).

It is contemplated that a released identifier nucleotide could be labeled before or after depolymerization using cross-linking chemistry well known in the art with commercially available reagents. For example, fluorescein isothiocyanate and rhodamine B isothiocyanate are both available from Aldrich Chemical Company (Milwaukee, Wisconsin). References to fluorescein isothiocyanate's use in labeling biological molecules include *Nature*, 193:167 (1962), *Methods Enzymol.* 26:28 (1972), *Anal. Biochem.*, 57:227 (1974), *Proc. Natl. Acad. Sci., U.S.*, 72:459 (1975).

It is contemplated that for many embodiments of the invention, it is useful to separate released fluorescent identifier nucleotides from those bound to an oligonucleotide, such as a probe. Thus, the separation techniques well known in

the art and discussed above are useful with such an embodiment, including HPLC fitted with a fluorescence detector. The enhanced sensitivity of fluorescence relative to other spectroscopic techniques can be  
5 used to increase the sensitivity of a detection or quantification process of the invention.

In an alternative embodiment wherein the analytical output is determined using fluorescence spectroscopy, an NADH detection system is used. In  
10 the NADH detection system, a combination of two enzymes, phosphoglycerate kinase and glyceraldehyde phosphate dehydrogenase, is used to catalyze the formation of NAD from NADH in the presence of ATP. Thus, this is in effect an ATP detection system, and  
15 much of the discussion herein relating to the detection of ATP with respect to the luciferase/luciferin system applies here. Because NADH is fluorescent whereas NAD is not, ATP is measured as a loss in fluorescence intensity.  
20 Examples of NADH based ATP assays are disclosed in United States Patent Nos. 4,735,897, 4,595,655, 4,446,231 and 4,743,561, and UK Patent Application GB 2,055,200, all of which are herein incorporated by reference.

25

#### **D. Absorbance Spectroscopic Analysis**

An absorbance spectrographic analysis step is contemplated to provide an analytical output, thereby provide for the determination of the presence  
30 or absence released identifier nucleotide, and indicate the presence or absence of said nucleic acid

target sequence. This embodiment contemplates the chromatographic separation of a reaction mixture that has been treated with a depolymerizing amount of an enzyme whose activity is to release one or more  
5 nucleotides from the 3'-terminus of a hybridized nucleic acid.

In an illustrative embodiment, a multiplexed assay for the presence of several different nucleic acid target sequences in a sample  
10 is analyzed by absorbance spectroscopy. Several labeled probes to various nucleic acid target sequences are added to a nucleic acid sample. The labels on the probes may be various nucleotide analogs, a different one for each probe. A  
15 depolymerizing enzyme is added, such as Klenow exo-, releasing the labeled nucleotides and other nucleotides from the 3'-termini of probes hybridized to target sequences when the 3' terminal nucleotide is matched.

20 The reaction solution is loaded onto a pre-equilibrated High Pressure Liquid Chromatography (HPLC) column and eluted under conditions that separate the nucleotide analogs from the natural nucleotides. Useful media for chromatographic  
25 separation of nucleotides, bases, and nucleosides include reverse phase media, such as a reverse phase C18 column or ODS-80T<sub>M</sub> or ODS-120T TSK-GEL by TosoHaas (Montgomeryville, Pennsylvania), anion exchange media, such as DEAE-25SW or SP-25W TSK-GEL by  
30 TosoHaas (Montgomeryville, Pennsylvania), or affinity media, such as Boronate-5PW TSK-GEL by TosoHaas

(Montgomeryville, Pennsylvania). Example 65 illustrates an embodiment of the present invention using HPLC.

The HPLC column is fitted with an  
5 absorbance detector to monitor the column effluent. Hence, "absorbance spectroscopy" for this type of analysis. Typical wavelengths for monitoring HPLC detection of nucleotides are 250 nm, 260 nm and 280 nm. Such separations of nucleotides and nucleotide  
10 analogs are well known in the art. Revich et al., *J. Chromatography*, 317:283-300 (1984), and Perrone & Brown, *J. Chromatography*, 317:301-310 (1984) provide examples of the HPLC separation of dNTPs.

Identification of the separated nucleotide  
15 analogs can be accomplished by comparison of the retention times (as monitored by absorbance of effluent at various times) of standards of the nucleotide analogs separated on the same HPLC column under the same conditions. Alternatively, the  
20 identity of the nucleotide analogs collected in separate fractions (as determined by continually monitoring the absorbance of the column effluent) can be determined by other standard analytical methods, such as nuclear magnetic resonance or atomic analysis  
25 (H,C,N).

In this illustrative example using  
depolymerization with Klenow exo-, the presence of a released identifier nucleotide from a particular  
probe indicates the presence of the target sequence  
30 that hybridize with that probe.

In an alternative embodiment, the released nucleotides from a depolymerization reaction mixture are separated on a gas chromatograph fitted with an absorbance detector to monitor column effluent.

5

#### **Probe-Mediated Specific Nucleic Acid Detection**

In yet another preferred embodiment, the probe-mediated specific nucleic acid detection method of the present invention can be used to simply  
10 identify or detect a nucleic acid of interest. For this method, a nucleic acid probe (e.g., DNA or RNA) is utilized which is substantially complementary to the target nucleic acid, which can be RNA or DNA. In a particularly preferred embodiment, the nucleic acid  
15 probe is entirely complementary to the exogenous target nucleic acid. The nucleic acid probe comprises single-stranded nucleic acid (e.g., DNA or RNA). The probe can be of varying lengths, preferably from about 10 to about 1000 bases, most  
20 preferably about 10 to 100 bases. Detection is carried out as described above. The nucleic acid probe-nucleic acid target/probe hybrid (complex) is exposed to conditions permitting depolymerization of the probe, which results in the production of XTPs.  
25 Detection of the nucleic acid of interest is characterized by a difference in the signal generated by the XTPs produced. Preferably, the XTPs are converted to ATP as described above and the ATP detected by a luciferase or NADH detection system.

30

Preferred conditions for depolymerization (depolymerization conditions) are described elsewhere

herein. The nucleotides are then detected. In some preferred embodiments, the nucleotides are converted to ATP equivalents as described hereinabove and in the Examples. In preferred embodiments, the ATP is  
5 detected by luciferase (luminescence spectroscopy) or NADH (fluorescence spectroscopy) detection systems.

As mentioned before, the determination of an appropriate nucleic acid target sequence useful for designing nucleic acid probes for use in a method  
10 of the invention is within the skill of the art.

Depolymerization reactions can also be used to interrogate the identity of a specific base in a nucleic acid. For example, the identity of single base point mutations, deletions, or insertions in a  
15 nucleic acid can be determined as follows.

In one embodiment, a nucleic acid probe is synthesized that is substantially complementary to a target nucleic acid containing or suspected of containing a point mutation. It will be recognized  
20 that various hybridization conditions can be used, so as to vary the stringency at which hybridization occurs. Thus, depending upon the system utilized, the complementarity of the probe can be varied. Depending on the length of the probe, the GC content,  
25 and the stringency of the hybridization conditions, the probe can have as many as 10 base mismatches with the target nucleic acid, and preferably less than 5 mismatches. Most preferably, the probe has only one base mismatch with the target nucleic acid or is  
30 completely complementary to the target nucleic acid.

The nucleic acid probe comprises single-stranded nucleic acid (e.g., DNA or RNA). The probe can be of varying lengths, preferably from about 10 to 100 bases, most preferably about 10 to 30 bases.

5 In particularly preferred embodiments, the probe is complementary to the target at all bases between an interrogation position and 3' end of the nucleic acid probe.

In preferred embodiments, the probe is

10 designed to have a predetermined nucleotide at an interrogation position. When the complementary probe base pairs or hybridizes to the target nucleic acid, the base at an interrogation position aligns with the base in the nucleic acid target whose identity is to

15 be determined under conditions such that base pairing can occur. It is contemplated that an interrogation position can be varied within the probe. For example, in some preferred embodiments, an interrogation position is preferably within 10 bases

20 of the 3' end of the nucleic acid probe. In still other preferred embodiments, an interrogation position is within 6 bases of the 3' end of the nucleic acid probe. In particularly preferred embodiments, an interrogation position is at the next

25 to last or last base at the 3' end of the nucleic acid probe.

In some preferred embodiments, four different probes of equal length are synthesized, each having a different nucleotide at an

30 interrogation position. Accordingly, it is contemplated that in some embodiments, a set of DNA

probes includes a first probe with a deoxyadenosine residue at an interrogation position, a second probe with a deoxythymidine residue at an interrogation position, a third probe with a deoxyguanosine residue at an interrogation position, and a fourth probe with a deoxycytosine residue at an interrogation position. Likewise, it is also contemplated that a set of RNA probes includes a first probe with an adenosine residue at an interrogation position, a second probe with a uridine residue at an interrogation position, a third probe with a guanosine residue at an interrogation position, and a fourth probe with a cytosine residue at an interrogation position.

In the next step of some embodiments, the probe or probes are hybridized to the exogenous target nucleic acid in separate reactions so that a probe nucleic acid-target nucleic acid complex is formed. It is contemplated that hybridization conditions can vary depending on the length and base composition of the probe.

In the probe-target nucleic acid complex, the nucleotide at an interrogation position is aligned with the specific base to be identified in the nucleic acid. In embodiments in which a set of probes is utilized, a different reaction is performed with each probe. In a multiplex embodiment, the set of probes can be used simultaneously. Because the probes differ at an interrogation position only one of the probes is complementary to the specific base in the target nucleic acid that is aligned with an interrogation position.



In the next step of some embodiments, the nucleic acid probe-target nucleic acid complexes are individually reacted under conditions permitting depolymerization of the hybridized probe. The preferred reaction conditions for depolymerization are described before and in the following Examples. The nucleotides are then detected.

In preferred embodiments, the reaction mixture also contains reagents necessary to catalyze the conversion of XTP to ATP equivalents as described in reaction 3 and in the following Examples. In some preferred embodiments, the nucleotides and/or ATP produced by the depolymerization reaction are then detected by either a luciferase or NADH detection system. Complementarity of the base at an interrogation position of the nucleic acid probe to the corresponding base in the nucleic acid target is characterized by detection of a signal generated from ATP following depolymerization.

In particularly preferred embodiments, the identity of the specific base is determined by comparing the amount of ATP produced in each reaction. Depolymerization of the probe proceeds from its 3' end. When the base at an interrogation position is not complementary to the specific base in the nucleic acid, very little or no ATP is produced, and thus no signal results. In alternative embodiments, this method can be practiced with from one to four probes. It is contemplated that utilizing multiple probes, (e.g., each with a different base at an interrogation position), may

prove unnecessary if a positive signal is produced (e.g., with the first probe tested).

#### **Assays Using Hairpin Structures**

5           Although it is preferred that the probes be constructed to be free of hairpin structures, assays in which hairpin structures are constructed are also useful. An embodiment of the invention, such as demonstrated in Example 20, contemplates use of a  
10 hairpin structure for determining the presence or absence of a nucleic acid target sequence in a nucleic acid sample with a probe that is hybridized to the exogenous target and then modified to be able to form a hairpin structure. This embodiment  
15 comprises the following steps.

          A treated sample is provided that contains a nucleic acid sample that may include an exogenous nucleic acid target sequence having an interrogation position. The target sequence, if present in the  
20 nucleic acid sample is hybridized with a nucleic acid probe. The probe is comprised of at least two sections. The first section contains the probe 3'-terminal about 10 to about 30 nucleotides. These nucleotides are complementary to the target strand  
25 sequence at positions beginning about 1 to about 30 nucleotides downstream of the interrogation position. The second section of the probe is located at the 5'-terminal region of the probe and contains about 10 to about 20 nucleotides of the target sequence. This  
30 same sequence, therefore, exists in both the target and the probe in the same 5' to 3' orientation. This

sequence spans the region in the target from the nucleotide at or just upstream (5') of the interrogation position, to the nucleotide just upstream to where the 3'-terminal nucleotide of the probe anneals to the target. An optional third section of the probe, from zero to about 50, preferably from zero to about 20, nucleotides in length and comprising a sequence that does not hybridize with either the first or second section, is located between the first and second sections of the probe.

The probe of the treated sample is extended in a template-dependent manner, by admixture with dNTPs and a template-dependent polymerase, at least through the interrogation position, thereby forming an extended probe/target hybrid that contains a sequence complementary to that at the interrogation position. In a preferred embodiment, the length of the probe extension is limited by omission from the extension reaction of a dNTP complementary to a nucleotide of the target sequence that is present upstream of the interrogation position and absent between the nucleotide complementary to the 3'-end of the interrogation position.

The extended probe/target hybrid is separated from any unreacted dNTPs; i.e., purified at least to the degree needed to use the extended probe strand to determine the presence or absence of the interrogation region in the sample or the identity of the base at the interrogation position. The extended probe/target hybrid is denatured to separate the

strands. The extended probe strand is permitted to form a hairpin structure.

A treated reaction mixture is formed by admixing the hairpin structure-containing composition with a depolymerizing amount of an enzyme whose activity is to release one or more nucleotides from the 3'-terminus of an extended probe hairpin structure. The reaction mixture is maintained under depolymerizing conditions for a time period sufficient for the depolymerizing enzyme to release 3'-terminus nucleotides, and then analyzed for the presence of released identifier nucleotides. The analytical output indicates the presence or absence of the exogenous nucleic acid target sequence. That analytical output can be determined as discussed elsewhere herein.

A still further embodiment of the invention, such as that termed REAPER™ and demonstrated in Example 21 and Fig. 2, also contemplates use of hairpin structures in determining the presence or absence of an exogenous nucleic acid target sequence, or a specific base within the target sequence, in a nucleic acid sample, and comprises the following steps. A treated sample is provided that contains a nucleic acid sample that may include an exogenous nucleic acid target sequence hybridized with a first nucleic acid probe strand (Fig. 2A).

The hybrid is termed the first hybrid. The first probe is comprised of at least two sections. The first section contains the probe 3'-terminal about 10 to about 30 nucleotides that are

complementary to the target nucleic acid sequence at a position beginning about 5 to about 30 nucleotides downstream of the target interrogation position. The second section of the first probe contains about 5 to about 30 nucleotides that are a repeat of the target sequence from the interrogation position to about 10 to about 30 nucleotides downstream of the interrogation position, and does not hybridize to the first section of the probe. That is, the second sequence is a repeat of the region in the exogenous target sequence from the interrogation position downstream to the position where the 3'-terminal nucleotide of the first probe aligns with the target. An optional third section of the probe, located between the first and second sections of the probe, is zero to about 50, preferably to about 20, nucleotides in length and comprises a sequence that does not hybridize to either the first or second section.

The first hybrid in the treated sample is extended at the 3'-end of the first probe, thereby extending the first probe past the interrogation position and forming an extended first hybrid (Fig. 2B) whose sequence includes an interrogation position and a sequence complementary to the exogenous target sequence and the exogenous target sequence interrogation position. The extended first hybrid is comprised of the original target nucleic acid and extended first probe. The extended first hybrid is then denatured in an aqueous composition to separate the two nucleic acid strands of the hybridized duplex

and form an aqueous solution containing a separated target nucleic acid and a separated extended first probe.

5 A second probe, that is about 10 to about 2000, more preferably about 10 to about 200, most preferably about 10 to about 30 nucleotides in length and is complementary to the extended first probe at a position beginning about 5 to about 2000, preferably about 5 to about 200, nucleotides downstream of the  
10 interrogation position in extended first probe, is annealed to the extended first probe, thereby forming the second hybrid (Fig. 2C). The second hybrid is extended at the 3'-end of the second probe until that extension reaches the 5'-end of the extended first  
15 probe, thereby forming a second extended hybrid (Fig. 2D) whose 3'-region includes an identifier nucleotide.

It is preferred that the polymerase enzyme utilized for an extension reaction be a template-  
20 dependent polymerase that is free of activity that adds a 3'-terminal deoxyadenosine in a template-nonspecific manner. Thus, it is preferred to use other than a polymerase such as *Taq* for a contemplated extension.

25 An aqueous composition of the extended second hybrid is denatured to separate the two nucleic acid strands; i.e., the extended second probe and the extended first probe. The aqueous composition so formed is cooled to form a "hairpin  
30 structure" from the separated extended second probe (Fig. 2E) when the target sequence is present in the

original nucleic acid sample. Thus, when the  
exogenous target sequence is present in the original  
nucleic acid sample, the 3'-terminal sequence of the  
second extended probe in the second extended hybrid  
5 hybridizes with the sequence of the second extended  
probe from a region comprising the interrogation  
position and nucleotides downstream from the  
interrogation position of second extended probe to  
the nucleotide position where the 3'-terminal  
10 nucleotide of the original (first-named) probe  
annealed to the original target.

A treated reaction mixture is formed by  
admixing the hairpin structure-containing composition  
with a depolymerizing amount of an enzyme whose  
15 activity is to release one or more nucleotides from  
the 3'-terminus of a nucleic acid hybrid. The  
reaction mixture is maintained under depolymerizing  
conditions for a time period sufficient to release  
3'-terminal region identifier nucleotides, and then  
20 analyzed for the presence of released identifier  
nucleotides. The analytical output indicates the  
presence or absence of the exogenous nucleic acid  
target sequence. Again, the analytical output can be  
determined by one of the several methods discussed  
25 elsewhere herein.

As was the case in the previous embodiment,  
dNTPs are utilized in the extension reactions. It is  
preferred that the hairpin structures be separated  
from the dNTPs prior to depolymerization to enhance  
30 the analysis for the identifier nucleotide.

## Kits

Other embodiments of the invention contemplate kits for determining the presence or absence of a predetermined exogenous nucleic acid target sequence in a nucleic acid sample. Such a kit comprises an enzyme whose activity is to release one or more nucleotides from the 3' terminus of a hybridized nucleic acid probe and at least one nucleic acid probe, said nucleic acid probe being complementary to the predetermined exogenous nucleic acid target sequence.

The kit optionally further comprises a nucleoside diphosphate kinase. Preferably, the nucleoside diphosphate kinase is that encoded by *Pyrococcus furiosus*. The kit optionally further comprises instructions for detecting the nucleic acid by depolymerization.

Preferably the enzyme whose activity is to release nucleotides in the kit is a template dependent polymerase that, in the presence of pyrophosphate ions, depolymerizes hybridized nucleic acids whose bases in the 3'-terminal region are matched with total complementarity. Alternatively, the enzyme whose activity is to release nucleotides in the kit exhibits a 3' to 5' exonuclease activity, depolymerizing hybridized nucleic acids having one or more mismatched bases at the 3' terminus of the hybridized probe.

It is to be understood that such a kit is useful for any of the methods of the present invention. The choice of particular components is



dependent upon the particular method the kit is designed to carry out. Additional components can be provided for detection of the analytical output, as measured by the release of identifier nucleotide, or  
5 by detection of the remaining probe after depolymerization. For example, luciferase assay reagent can be provided in the kits of the invention for detection of an identifier nucleotide released from the 3'-terminal region of a probe.

10 The instructions present in such a kit instruct the user on how to use the components of the kit to perform the various methods of the present invention. These instructions can include a description of the detection methods of the  
15 invention, including detection by luminescence spectroscopy, mass spectrometry, fluorescence spectroscopy, and absorbance spectroscopy.

In another embodiment, the invention contemplates a kit for determining the presence or  
20 absence of at least one predetermined exogenous nucleic acid target sequence in a nucleic acid sample comprising the following components: an enzyme whose activity, under depolymerizing conditions and in the presence of pyrophosphate is to release identifier  
25 nucleotide as a nucleoside triphosphate from hybridized nucleic acid probe; adenosine 5' diphosphate; pyrophosphate; a nucleoside diphosphate kinase; and at least one nucleic acid probe, wherein the nucleic acid probe is complementary to the  
30 predetermined exogenous nucleic acid target sequence.

Preferably, the enzyme whose activity in the presence of pyrophosphate is to release identifier nucleotides is the *Tne* triple mutant DNA polymerase, Klenow *exo-*, Klenow, T4 DNA polymerase, *Ath* DNA polymerase, *Taq* DNA polymerase and *Tvu* DNA polymerase, most preferably *Tne* triple mutant DNA polymerase, Klenow *exo-*, or *Tvu* DNA polymerase. In an alternative embodiment, a thermostable polymerase is preferred, wherein in the thermostable polymerase is preferably the *Tne* triple mutant DNA polymerase, T4 DNA polymerase, *Ath* DNA polymerase, *Taq* DNA polymerase and *Tvu* DNA polymerase, most preferably *Tne* triple mutant DNA polymerase, or *Tvu* DNA polymerase. Preferably, the nucleoside diphosphate kinase is that encoded by *Pyrococcus furiosus*. The kit optionally comprises instructions for use.

In another embodiment, the invention contemplates a kit for determining the presence or absence of a predetermined exogenous nucleic acid target sequence in a nucleic acid sample comprising an enzyme whose activity is to release one or more nucleotides from the 3' terminus of a hybridized nucleic acid probe and instructions for use. Such a kit optionally comprises a nucleoside diphosphate kinase. Preferably, the nucleoside diphosphate kinase is that encoded by *Pyrococcus furiosus*. The kit further optionally comprises a nucleic acid probe complementary to the predetermined exogenous nucleic acid target sequence.

In other embodiments of the present invention, nucleic acid detection test kits are

provided for performing a depolymerization method contemplated by this invention, and particularly a depolymerization detection method.

In one embodiment, the kit includes a  
5 vessel containing an enzyme capable of catalyzing pyrophosphorolysis, including, but not limited to *Taq* polymerase, *Tne* polymerase, *Tne* triple mutant polymerase, *Tth* polymerase, *Tvu* polymerase, *Ath* polymerase, T4 DNA polymerase, Klenow fragment,  
10 Klenow exo minus, *E. coli* DNA polymerase I, AMV reverse transcriptase, MMLV reverse transcriptase, or poly(A) polymerase, preferably a thermostable polymerase, most preferably *Tne* triple mutant polymerase or *Tvu* polymerase. In another embodiment,  
15 the kit contains a vessel that contains an exonuclease such as S1 nuclease, nuclease BAL 31, mung bean nuclease, exonuclease III and ribonuclease H.

Either of the above enzyme types is  
20 utilized in a contemplated method in a depolymerizing effective amount. That is, the enzyme is used in an amount that depolymerizes the hybridized probe to release an identifier nucleotide under depolymerizing conditions. This amount can vary with the enzyme  
25 used and also with the temperature at which depolymerization is carried out. An enzyme of a kit is typically present in an amount of about 0.1 to 100 U/reaction; in particularly preferred embodiments, the concentration is about 0.5 U/reaction. An amount  
30 of enzyme sufficient to carry out at least one assay, with its controls is provided.

As noted elsewhere, the preferred analytical output for determining the presence or absence of identifier nucleotide is luminescence caused by the reaction of ATP with luciferin in the presence of luciferase. A kit containing a pyrophosphorylation enzyme for use in DNA detection using luminescence also preferably includes a vessel containing NDPK and a vessel containing ADP. Similarly, a kit containing an exonuclease enzyme for use in DNA detection using luminescence also preferably includes a vessel containing PRPP synthetase and a vessel containing ADP. The NDPK or PRPP synthetase is provided in concentration of about 0.01 to 100 U/reaction, preferably about 0.1 to about 1.0 U/reaction.

Preferably, these reagents, and all of the reagents utilized in the kits discussed herein, are free of contaminating ATP and adenylate kinase. Some of the contaminants can be removed from the enzymes by dialysis treatment or by heat treatment.

Optionally, the kit contains vessels with reagents for amplification of dNTPs or NTP to ATP. Amplification reagents include, but are not limited to pyruvate kinase, adenylate kinase, NMPK, NDPK, AMP (e.g., as the amplification enzymes and substrate), and dCTP or AMP-CPP (e.g., as high-energy phosphate donors). In particularly preferred embodiments, the kit can be packaged in a single enclosure including instructions for performing the assay methods. In some embodiments, the reagents are provided in containers and are of a strength suitable for direct

use or use after dilution. In alternative preferred  
embodiments, a standard set can also be provided in  
order to permit quantification of results. In yet  
other preferred embodiments, test buffers for optimal  
5 enzyme activity are included.

In yet other embodiments, a contemplated  
kit comprises a nuclease, PRPP synthetase, PRPP,  
NDPK, and ADP together with luciferase and luciferin.  
In preferred embodiments, the nuclease is provided in  
10 a concentration of about 1 to 500 U/reaction; in  
particularly preferred embodiments at a concentration  
of about 20 U/reaction. In a particularly preferred  
embodiment, the PRPP synthetase is provided in  
concentration of about 0.01 U/reaction to 10  
15 U/reaction, preferably about 0.1 U/reaction. In some  
preferred embodiments, the kit includes all these  
reagents with luciferase and luciferin being provided  
as a single reagent solution.

In other preferred embodiments, these  
20 reagents include, but are not limited to, a high  
energy phosphate donor which cannot be utilized by  
luciferase, preferably dCTP, and AMP together with  
luciferase and luciferin. In alternative preferred  
embodiments, the kit includes all these reagents with  
25 luciferase and luciferin being provided in the same  
solution.

In still further embodiments of the present  
invention, the kits described above can contain a  
probe or probes for probe-mediated specific nucleic  
30 acid detection. In some embodiments, the kit  
contains at least one nucleic acid probe for a

nucleic acid target of interest. In other  
embodiments, the kits contain multiple probes, each  
of which contain a different base at an interrogation  
position or which are designed to interrogate  
5 different target DNA sequences.

The types of nucleic acid probes that can  
be included in the kits and their uses are described  
in greater detail herein.

10 Example 1: Reduction of Probe-alone Background  
Values for Probes Designed to  
Interrogate a Viral Sequence

In this example, the background light  
values from probe-alone reactions are reduced by  
15 alteration of reaction conditions. More  
specifically, the values from such background  
reactions are reduced by lowering the Klenow exo-  
level in the reactions. In addition, the probes are  
used to assay the relative probe signal strength  
20 values for probes that hybridize to the same DNA  
strand versus probes that hybridize to different  
strands but that interrogate the same nucleotide  
polymorphism site.

Oligonucleotides CV11 (SEQ ID NO:1) and  
25 CV12 (SEQ ID NO:2) are a pair of single-stranded DNAs  
that can hybridize together to produce a segment of  
the genome of cytomegalovirus (CMV) in a form  
sensitive to the drug gancyclovir. Oligonucleotides  
CV13 (SEQ ID NO:3) and CV14 (SEQ ID NO:4) are a pair  
30 of single-stranded DNAs that can hybridize together  
to produce the same segment of the CMV genome, but

differ from CV11 and CV12 in that they contain a SNP that represents a form of the virus resistant to the drug gancyclovir.

Probe oligonucleotide CV15 (SEQ ID NO:5)  
5 can hybridize with exact homology to a segment of CV12. Probe oligonucleotide CV16 (SEQ ID NO:6) is identical to CV15 except that it contains a one base change from the CV15 sequence at the site of the SNP that confers drug resistance to the virus. Probe  
10 oligonucleotide CV17 (SEQ ID NO:7) can hybridize with exact homology to CV11. Probe oligonucleotide CV18 (SEQ ID NO:8) is identical to CV17 except that it contains a one base change from the CV17 sequence at the site of the SNP that confers drug resistance to  
15 the virus.

The oligonucleotides above were dissolved in water at a concentration of 1 mg/mL and the following solutions were assembled.

Solution	Oligonucleotide	Water
#1	--	20 µL
#2	CV15, 1 µL	19 µL
#3	CV16, 1 µL	19 µL
#4	CV17, 1 µL	19 µL
#5	CV18, 1 µL	19 µL

20

These solutions were heated at 95°C for 5 minutes, then cooled at room temperature for 10 minutes. The following master mix was assembled and mixed.

Component	Amount
10X DNA Pol Buffer (Promega, M195A)	200 $\mu$ L
Klenow exo- (1U/ $\mu$ L) (Promega M218B)	12.5 $\mu$ L
40 mM Sodium Pyrophosphate (Promega C350B)	25 $\mu$ L
NDPK (1U/ $\mu$ L)	10 $\mu$ L
10uM ADP (Sigma A5285)	20 $\mu$ L
Water	732.5 $\mu$ L

Twenty microliters of this solution were added to solutions 1-5 above after they had cooled, and then the resulting mixtures were heated at 37°C for 15 minutes. After this incubation, 4  $\mu$ L of each solution were added to 100  $\mu$ L of L/L reagent (Promega F202A) and the light production of the resulting solution was measured immediately using a Turner® TD 20/20 luminometer. The following results were obtained.

Solution sampled	Relative light units
#1	13.07
#2	14.98
#3	14.27
#4	28.25
#5	583.70



These results demonstrate that probes CV15-  
CV17 provide relatively low probe-alone light signals  
at 0.25U Klenow exo- per reaction but that probe  
CV18-alone provides a very high relative light  
5 signal. The sequence of the CV18 probe can form a  
hairpin structure such that the terminal 3' bases  
hybridize to the sequence 5'TCGTGC 3' further towards  
the 5' end of the segment. Although probe CV17 could  
form the same structure, the terminal 3' base of the  
10 resulting structure would have a mispaired base.

These data exemplify one of the guiding  
principles of appropriate probe design for this  
system: the probes should not be predicted to form  
stable hairpin structure and, in particular, should  
15 not be predicted to give such a structure with the 3'  
end of the probe producing a structure that forms a  
blunt end or 5' overhang in the fragment as they may  
act as a substrate for the depolymerizing enzyme. In  
addition, the probes used should not be predicted to  
20 form probe dimer structures with either blunt ends or  
5' overhanging ends because such probes can produce  
high probe-alone signals in the system and might make  
them unacceptable for use.

Due to their low background, probes CV15-  
25 CV17 were then selected for further study. Equal  
volumes of oligonucleotides CV11 and CV12 were  
annealed together, as described earlier in this  
example, as were CV13 and 14. The annealed solutions  
of CV11 and CV12, and CV13 and CV14 were labeled  
30 CV11+12 and CV13+14, respectively. The following  
solutions were assembled.

Solution	CV15	CV16	CV17	CV11+12	CV13+14	CV(11+12) + (13+14) Heterozyg Template	Water
#1	--	--	--	--	--	--	20 $\mu$ L
#2	1 $\mu$ L	--	--	--	--	--	19 $\mu$ L
#3	--	1 $\mu$ L	--	--	--	--	19 $\mu$ L
#4	--	--	1 $\mu$ L	--	--	--	19 $\mu$ L
#5	--	--	--	1 $\mu$ L	--	--	19 $\mu$ L
#6	--	--	--	--	--	1 $\mu$ L	19 $\mu$ L
#7	--	--	--	--	1 $\mu$ L	--	19 $\mu$ L
#8	1 $\mu$ L	--	--	1 $\mu$ L	--	--	18 $\mu$ L
#9	--	1 $\mu$ L	--	1 $\mu$ L	--	--	18 $\mu$ L
#10	1 $\mu$ L	--	--	--	--	1 $\mu$ L	18 $\mu$ L
#11	--	1 $\mu$ L	--	--	--	1 $\mu$ L	18 $\mu$ L
#12	1 $\mu$ L	--	--	--	1 $\mu$ L	--	18 $\mu$ L
#13	--	1 $\mu$ L	--	--	1 $\mu$ L	--	18 $\mu$ L
#14	1 $\mu$ L	--	--	1 $\mu$ L	--	--	18 $\mu$ L
#15	--	--	1 $\mu$ L	1 $\mu$ L	--	--	18 $\mu$ L
#16	1 $\mu$ L	--	--	--	--	1 $\mu$ L	18 $\mu$ L
#17	--	--	1 $\mu$ L	--	--	1 $\mu$ L	18 $\mu$ L
#18	1 $\mu$ L	--	--	--	1 $\mu$ L	--	18 $\mu$ L
#19	--	--	1 $\mu$ L	--	1 $\mu$ L	--	18 $\mu$ L

These solutions were heated at 95°C for 5 minutes and then permitted to cool for 10 minutes at room temperature. A master mix solution was assembled as in described in this Example, containing Klenow exo- at a final concentration of 0.25U/20  $\mu$ L. After solutions 1-19 had cooled, 20  $\mu$ L of the master mix solution were added and the resulting solution heated at 37°C for 15 minutes. After this incubation,

duplicate 4  $\mu$ L samples of solutions 2-19 and a single sample of solution 1 were taken, added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light production of the mixture measured immediately using a Turner®  
5 TD 20/20 luminometer. The following results were obtained.

Relative light units		
Solution	Reading 1	Reading 2
#1	10.53	--
#2	11.35	12.16
#3	10.79	12.75
#4	17.70	16.76
#5	12.78	11.12
#6	11.36	11.48
#7	12.38	12.16
#8	348.3	369.3
#9	73.11	74.48
#10	289.5	283.6
#11	509.8	364.0
#12	120.2	108.6
#13	785.4	595.7
#14	764.3	763.3
#15	77.25	73.22
#16	530.9	541.2
#17	476.1	419.6
#18	339.4	262.7
#19	943.2	964.0

The net relative light values for the data above were calculated as follows. The ratios reported in this example were determined by first averaging the results from matching samples, then  
5 determining the net light production from the matching and mismatching samples and dividing the net light production from the matching reaction by that seen in the mismatch reaction. The net light production was determined by subtracting the  
10 estimated light contribution from the probes and template present in the reactions from the total light produced. The light production from the template reaction was considered to be the total of that contributed from the template specifically and  
15 that contributed by contaminating ATP from various components. The net increase from the probes alone was calculated by subtracting the average "No DNA" values from the probe values since this subtracts the contributions from contaminating ATP from the probe  
20 values. Thus, the formula used to determine the net light production from the reactions was:

$$\begin{aligned} \text{Net Light} = & \text{Total light} - [(\text{target alone}) \\ & + (\text{probe alone} - \text{No DNA})] \end{aligned}$$

These values were used to determine the  
25 signal ratio by dividing the signal from the "C" allele probe by the signal from the "T" allele probe. The results of these calculations are presented in the tables below, wherein "WT" indicates the wild  
type genotype.

Probes Interrogate the Same DNA Strand				Probes Interrogate Different DNA Strands			
Template Genotype				Template Genotype			
Probe	C/C	C/T	T/T		C/C	C/T	T/T
WT Probe (CV15)	345.5	274.0	100.8	WT Probe (CV15)	745.1	518.0	282.1
Mutant Probe (CV16)	60.5	424.3	677	Mutant Probe (CV17)	61.9	435.0	940
Ratio	5.7	1.5	0.15	Ratio	12	1.2	0.33

These data demonstrate that, for this particular SNP, probes that detect the polymorphism that bind to different strands provide the signal ratio closest to 1.0 when both nucleic acid targets are present in the reaction (as occurs for samples heterozygous for a particular allele). However either set of probes give clearly different signals depending upon the genotype of the sample DNA.

CV11

5' CGCTTCTACCACGAATGCTCGCAGACCATGCTGCACGAATACGTCAGAAAG  
AACGTGGAGCGTCTGTTGGAGCT 3' SEQ ID NO:1

CV12

5' CCAACAGACGCTCCACGTTCTTTCTGACGTATTCGTGCAGCATGGTCTGCG  
AGCATTCGTGGTAGAAGCGAGCT 3' SEQ ID NO:2

CV13

5' CGCTTCTACCACGAATGCTCGCAGATCATGCTGCACGAATACGTCAGAAA  
GAACGTGGAGCGTCTGTTGGAGCT 3' SEQ ID NO:3

CV14

5' CCAACAGACGCTCCACGTTCTTTCTGACGTATTCGTGCAGCATGATCTGCG  
AGCATTCGTGGTAGAAGCGAGCT 3' SEQ ID NO:4

5 CV15 5' CTACCACGAATGCTCGCAGAC 3' SEQ ID NO:5

CV16 5' CTACCACGAATGCTCGCAGAT 3' SEQ ID NO:6

CV17 5' TGACGTATTCGTGCAGCATGG 3' SEQ ID NO:7

10

CV18 5' TGACGTATTCGTGCAGCATGA 3' SEQ ID NO:8

15 Example 2: Detection of DNA Sequences in  
the Genome of *Listeria* Species

This example provides an assay for the presence of DNA sequences present in the genome of *Listeria* in a gene known as the *iap* gene.

20 Oligonucleotides LM1 (SEQ ID NO:9) and LM2 (SEQ ID NO:10) encode a segment of the *iap* gene and are exactly complementary to each other. Oligonucleotide probe LM3 (SEQ ID NO:11) was designed to hybridize exactly with a region of target LM2, and probe LM4 (SEQ ID NO:12) was designed to hybridize exactly to  
25 target LM1.

Oligonucleotides LM1-LM4 were dissolved in TE buffer (10 mM Tris, 1 mM EDTA, pH8.0) at a concentration of 500 µg/mL and then were diluted 25-fold in TE buffer to obtain solutions at a DNA  
30 concentration of 20 ng/µL. The following solutions were assembled.

Solution	Oligonucleotides	1X TE Buffer
#1	LM1, 10 $\mu$ L	10 $\mu$ L
#2	LM2, 10 $\mu$ L	10 $\mu$ L
#3	LM3, 10 $\mu$ L	10 $\mu$ L
#4	LM4, 10 $\mu$ L	10 $\mu$ L
#5	LM1, 10 $\mu$ L; LM3, 10 $\mu$ L	--
#6	LM1, 10 $\mu$ L; LM4, 10 $\mu$ L	--
#7	LM2, 10 $\mu$ L; LM3, 10 $\mu$ L	--
#8	LM2, 10 $\mu$ L; LM4, 10 $\mu$ L	--
#9	--	20 $\mu$ L

5      These solutions were heated at 95°C for 3 minutes, then permitted to cool at room temperature for 15 minutes.

The following master mix was assembled.

Component	Volume/reaction
Nanopure water (Promega AA399)	12.75 $\mu$ L
10X DNA Polymerase Buffer (Promega M195A)	2 $\mu$ L
40 mM Sodium Pyrophosphate (Promega C113)	0.25 $\mu$ L
ADP, 2 $\mu$ M*	1 $\mu$ L
NDPK, 0.1U/ $\mu$ L**	1 $\mu$ L
Klenow Exo- 10U/ $\mu$ L (Promega M218)	1 $\mu$ L

\* Made by dissolving Sigma A5285 in water. \*\* Made by dissolving Sigma N0379 in water.

After solutions 1-9 had cooled, 2  $\mu$ L samples of the solution were added to 18  $\mu$ L of the master mix, in triplicate, the resulting solutions were mixed and incubated at 37°C for 15 minutes.

5 After this incubation, the tubes were placed on ice. Once all the incubations were on ice, 20  $\mu$ L of the contents of the tubes were added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light production of the resulting reaction was measured immediately using  
10 a Turner® TD 20/20 luminometer. The following data were obtained.

Relative light units						
Solution	Target	Probe	Reading 1	Reading 2	Reading 3	Avg.
#1	LM1	--	70.3	69.7	69.0	69.7
#2	LM2	--	39.6	40.8	45.3	41.9
#3	--	LM3	12.2	12.4	13.2	12.6
#4	--	LM4	16.9	17.3	17.4	17.2
#5	LM1	LM3	57.7	76.5	72.7	69.0
#6	LM1	LM4	1814	1815	1761	1797
#7	LM2	LM3	56.72	61.1	57.59	58.5
#8	LM2	LM4	67.5	72.4	79.3	73.1

These data show that LM4 produces a strong  
15 signal in the reaction with LM1 and thus can be used to detect this DNA sequence.

Oligonucleotides LM1 and LM2 were diluted to 2 ng/ $\mu$ L in 1X TE buffer. These materials were also used to create the following solutions in  
20 triplicate.



Solution	LM1	LM2	LM3	LM4	1X TE
#1	5 $\mu$ L	5 $\mu$ L	--	--	10 $\mu$ L
#2	5 $\mu$ L	5 $\mu$ L	10 $\mu$ L	--	--
#3	5 $\mu$ L	5 $\mu$ L	--	10 $\mu$ L	--

These solutions were heated to 95°C for 10 minutes, then permitted to cool for 15 minutes at room temperature.

5 A master mix was made as described earlier in this example. After cooling at room temperature, 2  $\mu$ L of each solution were added to an 18  $\mu$ L sample of this master mix, and the resulting solutions were incubated at 37°C for 15 minutes. After this  
10 incubation, 2  $\mu$ L of the solution were added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light produced was immediately read using a Turner® TD 20/20 luminometer. The following results were obtained.

15

Relative light units					
Solution	Reading 1	Reading 2	Reading 3	Avg.	NLU*
#1	754.4	727.8	752.7	745.0	--
#2	857.4	801.0	852.3	836.9	91.9
#3	1185	1211	1192	1196	451

\* Net light units (NLU) were calculated by subtracting the no probe reaction average (#1) from the specific probe reaction values.

20

With both DNA template strands present, both probes provide signals above background.

The sequences used were as follows:

LM1

5' GAAGTAAAACAAACTACACAAGCAACTACACCTGCGCCTAAAG  
5 TAGCAGAAACGAAAGAAACTCCAGTAG 3' SEQ ID NO:9

LM2

5' CTACTGGAGTTTCTTTTCGTTTCTGCTACTTTAGGCGCAGGT  
GTAGTTGCTTGTGTAGTTTGTTTTACTTC 3' SEQ ID NO:10  
10

LM3 5' GCAACTACACCTGCGCCTAAAGTAGCAGAA 3'  
SEQ ID NO:11

LM4 5' TTCTGCTACTTTAGGCGCAGGTGTAGTTCG 3'  
15 SEQ ID NO:12

Example 3: Detection of Segments of  
the *Listeria hyl* Gene

In this example, a method is described for  
20 the detection of a segment of the *hyl* gene from  
*Listeria monocytogenes*.

Oligonucleotides LM5 (SEQ ID NO:13) and LM6  
(SEQ ID NO:14) anneal exactly to create a region of  
the *hyl* gene. LM7 (SEQ ID NO:15) and LM8 (SEQ ID  
25 NO:16) oligonucleotides are used as interrogation  
probes with LM7 completely complementary to LM6 and  
LM8 completely complementary to LM5.

Oligonucleotides LM5-8 were dissolved in 1X TE buffer  
at a concentration of 500 µg/mL and then were diluted  
30 25 fold in TE buffer to obtain solutions at a DNA

concentration of 20 ng/μL. The following solutions were assembled.

Solution	Oligonucleotides	1X TE Buffer
#1	LM5, 10 μL	10 μL
#2	LM6, 10 μL	10 μL
#3	LM7, 10 μL	10 μL
#4	LM8, 10 μL	10 μL
#5	LM5, 10 μL; LM7, 10 μL	--
#6	LM5, 10 μL; LM8, 10 μL	--
#7	LM6, 10 μL; LM7, 10 μL	--
#8	LM6, 10 μL; LM8, 10 μL	--
#9	--	20 μL

5                    These solutions were heated at 95°C for 3 minutes, then permitted to cool at room temperature for 15 minutes.

The following master mix was assembled.

Volume/reaction	
Nanopure water(Promega AA399)	12.75 μL
10X DNA Polymerase Buffer(Promega M195)	2 μL
40 mM Sodium Pyrophosphate(Promega C113)	0.25 μL
ADP, 2 μM*	1 μL
NDPK, 0.1U/μL**	1 μL
Klenow Exo- 10U/μL (Promega M128)	1 μL

10                    \* Made by dissolving Sigma A5285 in water.    \*\* Made by dissolving Sigma N0379 in water.

After solutions 1-9 had cooled, triplicate  
2  $\mu$ L samples of the solution were added to 18  $\mu$ L  
master mix and the resulting solutions were mixed and  
incubated at 37°C for 15 minutes. After this  
5 incubation, the tubes were placed on ice. Once all  
the incubations were on ice, 20  $\mu$ L of the contents of  
the tubes were added to 100  $\mu$ L of L/L reagent  
(Promega F202A) and the light production of the  
resulting reaction was measured immediately using a  
10 Turner® TD 20/20 luminometer. The following data  
were obtained.

Relative light units					
Solution	Reading 1	Reading 2	Reading 3	Avg.	Net Ave
#1	28.53	29.62	30.0	29.41	---
#2	81.30	75.12	74.68	77.03	---
#3	19.88	13.12	12.80	15.26	---
#4	1326	1273	1216	1271	---
#5	37.24	36.40	36.77	36.80	3.78
#6	2582	2336	2169	2362	1089
#7	90.74	90.83	90.64	90.64	9.97
#8	1596	1671	1787	1684	347.6
#9	12.33	11.16	11.48	11.66	---

15 The above data indicate that at least  
oligonucleotide LM8 can be used to detect the target  
gene sequence represented in LM6.

Oligonucleotides LM5 and LM6 were diluted  
to 2 ng/ $\mu$ L in 1X TE buffer (10 mM Tris, 1 mM EDTA, pH

8.0). These materials were also used to create the following solutions in triplicate.

Solution	LM5	LM6	LM7	LM8	1X TE
#1	5 $\mu$ L	5 $\mu$ L	--	--	10 $\mu$ L
#2	5 $\mu$ L	5 $\mu$ L	10 $\mu$ L	--	--
#3	5 $\mu$ L	5 $\mu$ L	--	10 $\mu$ L	--

5                    These solutions were heated to 95°C for 10 minutes, and then cooled for 15 minutes at room temperature.

                  Then 2  $\mu$ L of the solutions were added to triplicate 18  $\mu$ L samples of the master mix and then  
10                   the resulting solutions were incubated at 37°C for 15 minutes. After this incubation, 2  $\mu$ L of the solution were added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light produced was immediately read using a Turner® TD 20/20 luminometer. The following results  
15                   were obtained.

Relative light units					
Solution	Reading 1	Reading 2	Reading 3	Avg.	NLU *
#1	442.5	431.8	432.2	435.5	--
#2	576.1	544.6	580.1	566.9	115.7
#3	1779	1837	1908	1841	1405

\*Net light units (NLU) determined by subtraction of probe alone values (see table above) and solution #1 values from the average light units measured.

20

                  These results demonstrate that specific detection of the segment of the *hyl* gene sequence from *Listeria* can be performed using the components

described above. Because this gene sequence is specific for *Listeria*, this indicates that the components can be used for specific detection of *Listeria* DNA.

5

LM5

5' CATCGACGGCAACCTCGGAGACTTACGAGATATTTTGAAAAA  
GGCGCTACTTTTAATCGAGAAACACCA 3' SEQ ID NO:13

10

LM6

5' TGGTGTTTCTCGATTAAAAGTAGCGCCTTTTTTCAAATATCT  
CGTAAGTCTCCGAGGTTGCCGTCGATG 3' SEQ ID NO:14

LM7 5' CTCGGAGACTTACGAGATATTTTGAAAAA 3' SEQ ID NO:15

15

LM8 5' TTTTTTCAAATATCTCGTAAGTCTCCGAG 3' SEQ ID NO:16

20

Example 4: Detection of a DNA Sequence  
from Salmonella

In this example, a method for detection of a gene sequence from *Salmonella* is provided.

25

Oligonucleotides ST1 (SEQ ID NO:17), ST2 (SEQ ID NO:18), ST3 (SEQ ID NO:19), and ST4 (SEQ ID NO:20) were dissolved in 1X TE buffer to 500 µg/µL and then were diluted 25 fold in 1X TE buffer to obtain solutions at a DNA concentration of 20 ng/µL. The following solutions were prepared.

Solution	Oligonucleotides	1X TE Buffer
----------	------------------	--------------

Solution	Oligonucleotides	1X TE Buffer
#1	ST1, 10 $\mu$ L	10 $\mu$ L
#2	ST2, 10 $\mu$ L	10 $\mu$ L
#3	ST3, 10 $\mu$ L	10 $\mu$ L
#4	ST4, 10 $\mu$ L	10 $\mu$ L
#5	ST1, 10 $\mu$ L; ST3, 10 $\mu$ L	--
#6	ST1, 10 $\mu$ L; ST4, 10 $\mu$ L	--
#7	ST2, 10 $\mu$ L; ST3, 10 $\mu$ L	--
#8	ST2, 10 $\mu$ L; ST4, 10 $\mu$ L	--
#9	--	20 $\mu$ L

These solutions were heated at 95°C for 3 minutes, then permitted to cool at room temperature for 15 minutes.

- 5 The following master mix was assembled.

Component	Volume/reaction
Nanopure water (Promega AA399)	12.75 $\mu$ L
10X DNA Polymerase Buffer (Promega M195)	2 $\mu$ L
40mM Sodium Pyrophosphate (Promega C113)	0.25 $\mu$ L
ADP, 2 $\mu$ M*	1 $\mu$ L
NDPK, 0.1U/ $\mu$ L**	1 $\mu$ L
Klenow Exo- 10U/ $\mu$ L (Promega M128)	1 $\mu$ L

\* Made by dissolving Sigma A5285 in water. \*\* Made by dissolving Sigma N0379 in water.

- 10 After solutions 1-9 had cooled, three 2  $\mu$ L samples of the solution were added to 18  $\mu$ L of the master mix and the resulting solution was mixed and

incubated at 37°C for 15 minutes. After this incubation, the tubes were placed on ice. Once all the incubations were on ice, 20 µL of the contents of the tubes were added to 100 µL of L/L reagent, and  
5 the light production of the resulting reaction was measured immediately using a Turner® TD 20/20 luminometer. The following data were obtained.

Relative light units					
Solution	Reading 1	Reading 2	Reading 3	Avg.	Net Avg.
#1	18.28	18.27	17.97	18.17	---
#2	231.9	211.4	226.3	223.2	---
#3	11.58	12.56	11.34	11.83	---
#4	14.00	14.48	14.88	14.45	---
#5	21.31	21.20	19.44	20.65	2.18
#6	3003	2943	2918	2955	2933
#7	2780	2782	2641	2734	2510
#8	256.4	269.9	271.1	265.8	39.67
#9	11.63	11.39	11.56	11.52	---

10                    These data indicate that both  
oligonucleotide probes ST3 and ST4 can give a very  
strong specific light signals with single strand  
target DNA sequence from *Salmonella*.

                    Oligonucleotides ST1 and ST2 were diluted  
15 to 2 ng/µL in 1X TE buffer (10mM Tris, 1mM EDTA, pH  
8.0). These materials were also used to create the  
following solutions in triplicate.

Solution	ST1	ST2	ST3	ST4	1X TE
#1	5 µL	5 µL	--	--	10µl



#2	5 µL	5 µL	10 µL	--	--
#3	5 µL	5 µL	--	10 µL	--

These solutions were heated to 95°C for 10 minutes, then permitted to cool for 15 minutes at room temperature.

5           A master mix was made as described earlier in this example. After cooling at room temperature, 2 µL of each solution were added to an 18 µL sample of this master mix, and then the resulting solutions were incubated at 37°C for 15 minutes. After this  
10 incubation, 2 µl of the solution were added to 100 µL of L/L reagent and the light produced was immediately read using a Turner® TD 20/20 luminometer. The following results were obtained.

Relative light units					
Solution	Reading 1	Reading 2	Reading 3	Avg.	NLU *
#1	692.5	728.9	678.3	699.9	--
#2	2448	2389	2311	2382	1683
#3	1742	1778	1738	1752	1053

15           \*Net light units (NLU) were determined by subtraction of probe alone values (see table above) and solution #1 values from the average light units measured.

20           These data demonstrate that oligonucleotide probes ST3 and ST4 provide specific detection of the DNA target sequence from *Salmonella* even if both DNA strands are present.

25           Sequences used were as follows:

ST1

5' TTTAATTCGGAGCCTGTGTAATGAAAGAAATCACCGTCACTG

AACCTGCCTTTGTCACC 3'

SEQ ID NO:17

5 ST2

5' GGTGACAAAGGCAGGTTTCAGTGACGGTGATTTCTTTCATTACACAGGCT

CCGGAATTAAA 3'

SEQ ID NO:18

ST3 5' TGTGTAATGAAAGAAATCACCGTCACTGAA 3'

10

SEQ ID NO:19

ST4 5' TTCAGTGACGGTGATTTCTTTCATTACACA 3'

SEQ ID NO:20

15 Example 5: Detection of a Specific  
Message by Use of a DNA Probe Exactly  
Matching the Message Sequence and Lack  
of a Signal when the DNA Probe is  
Mismatched at Its 3' End

20 In this Example, a luciferase light signal  
is generated from pyrophosphorylation of a DNA probe  
that complements the sequence of a target RNA  
species. In addition, evidence is presented to  
demonstrate that this signal is not generated if the  
25 3'-terminal base of the probe does not complement the  
RNA base in the target sequence. These data  
demonstrate that probe pyrophosphorylation can be  
used to detect the presence of specific target RNA  
sequences and that mutations at specific bases in the  
30 target sequence can be detected by use of probes that  
should match the base but that do not give a signal  
with the message.

A master mix was assembled that contained:

	Capped Kanamycin RNA (0.62 mg/mL)	1.25 $\mu$ L
	5X MMLV Reaction Buffer	50 $\mu$ L
	40 mM Sodium Pyrophosphate	2.5 $\mu$ L
5	10 $\mu$ M ADP	2.5 $\mu$ L
	NDPK (1 U/ $\mu$ L)	5 $\mu$ L
	MMLV-RT (200 U/ $\mu$ L) (Promega, M1701)	12.5 $\mu$ L
	Nanopure water	163.75 $\mu$ L

10 Probes one through four were dissolved at a concentration of 1mg/mL in 1X TE buffer.

Probe 1 (SEQ ID NO:21) was designed to exactly complement a segment of the coding region of the Kanamycin RNA. Probe 2 (SEQ ID NO:22), Probe 3 (SEQ ID NO:23) and Probe 4 (SEQ ID NO:24) were  
15 designed to match the sequence of Probe 1 except that the 3'-terminal base of the probe was altered to one of each of the other three DNA bases at this position.

20 Nineteen microliters of the master reaction mix were placed in 10 labeled 0.5 mL microfuge tubes and the following additions were made to the tubes:  
Tubes 1 and 2, 1  $\mu$ L 1X TE buffer; Tubes 3 and 4, 1  $\mu$ L Probe 1; Tubes 5 and 6, 1  $\mu$ L Probe 2; Tubes 7 and 8,  
25 1  $\mu$ L Probe 3; and Tubes 9 and 10, 1  $\mu$ L Probe 4. The 10 0.5 mL microfuge tubes were incubated at 37°C for 20 minutes to hybridize and form treated samples.  
Thereafter, 2  $\mu$ L of the contents of the tubes were added to 100  $\mu$ L L/L reagent (Promega, F202A) and the

light output of the reagent was measured using a luminometer. The following data were collected.

Solution	Relative Light Units
1	3.989
2	3.458
3	49.95
4	52.24
5	3.779
6	3.727
7	4.394
8	4.163
9	7.879
10	7.811

- 5 These data show that MMLV-RT is able to pyrophosphorylate a DNA probe that hybridized to a target RNA sequence and that the free nucleoside triphosphates that are formed are converted to ATP equivalents that can be measured using luciferase.
- 10 In addition, the data show that this signal is either absent or much weaker (solutions 1,2,5,6,7,8,9,10) when a probe with a 3' mismatch to the expected base is used in the reaction (compare to tubes 3 and 4).

Probe 1	SEQ ID NO:21	5'GCAACGCTACCTTTGCCATGTTTC 3'
Probe 2	SEQ ID NO:22	5'GCAACGCTACCTTTGCCATGTTTG 3'
Probe 3	SEQ ID NO:23	5'GCAACGCTACCTTTGCCATGTTTA 3'
Probe 4	SEQ ID NO:24	5'GCAACGCTACCTTTGCCATGTTTT 3'

Example 6:      Detection of a Specific RNA:  
                 Globin mRNA

5

In this Example, the light signal produced from pyrophosphorylation of DNA probes that are complementary to two regions of globin mRNA is compared to the signals from two DNA probes that are the exact sequence of the same regions. Once again, probes that totally complement the target RNA are shown to give a signal above background, whereas those that do not complement the target RNA give little or no signal.

15                Probe 5 (SEQ ID NO:25), Probe 6 (SEQ ID NO:26), Probe 7 (SEQ ID NO:27), and Probe 8 (SEQ ID NO:28) were diluted to a concentration of 0.5 mg/mL in 1X TE buffer (10 mM Tris, 1 mM EDTA). Purified globin mRNA (Gibco BRL, 18103-028) as target was dissolved in 1X TE buffer (10 mM Tris, 1 mM EDTA) to a concentration of 20 ng/ $\mu$ L.

Hybridization solutions were assembled as follows:

25    Solution 1:      10  $\mu$ L Probe 5 and 10  $\mu$ L Globin mRNA  
      Solution 2:      10  $\mu$ L Probe 6 and 10  $\mu$ L Globin mRNA  
      Solution 3:      10  $\mu$ L Probe 7 and 10  $\mu$ L Globin mRNA  
      Solution 4:      10  $\mu$ L Probe 8 and 10  $\mu$ L Globin mRNA  
      Solution 5:      10  $\mu$ L Probe 5 and 10  $\mu$ L 1X TE buffer  
30    Solution 6:      10  $\mu$ L Probe 6 and 10  $\mu$ L 1X TE buffer  
      Solution 7:      10  $\mu$ L Probe 7 and 10  $\mu$ L 1X TE buffer

Solution 8: 10  $\mu$ L Probe 8 and 10  $\mu$ L 1X TE buffer

Solution 9: 10  $\mu$ L 1X TE buffer, 10  $\mu$ L Globin mRNA

5 These solutions were assembled in 0.5 mL tubes, heated to 50°C for 15 minutes and permitted to cool to room temperature for 15 minutes.

The following master reaction mixture was assembled:

Nanopure water	346.5 $\mu$ L
MMLV-RT 5X Reaction Buffer (Promega M195A)	132 $\mu$ L
Sodium pyrophosphate (Promega M531)	16.5 $\mu$ L
NDPK (1 U/ $\mu$ L)	33 $\mu$ L
ADP (2 $\mu$ M)	33 $\mu$ L
MMLV-RT (adjusted to 100 U/ $\mu$ L) (Promega, M1701)	33 $\mu$ L

10

The solution above was mixed and 18  $\mu$ L placed into 27 tubes. Three two-microliter samples of each of the hybridization solutions above were added in three of the tubes containing the master  
15 reaction mix and the tubes were then incubated at 37°C for 15 minutes and permitted to cool to room temperature to hybridize and form treated samples. The contents of the tubes were then added to 100  $\mu$ L of L/L reagent and the light production of the  
20 resulting reaction was measured immediately using a luminometer (Turner® TD20/20). The following results were obtained:

Hybridization Solution	Light Values			Average
Probe 5 + RNA	6.555	6.303	6.187	6.348
Probe 5 + TE Buffer	6.335	5.923	6.046	6.101
Probe 6 + RNA	137.8	128.5	169.2	145.2
Probe 6 + TE Buffer	10.24	9.429	9.858	9.842
Probe 7 + RNA	6.235	6.763	6.375	6.458
Probe 7 + TE Buffer	6.436	6.545	6.138	6.388
Probe 8 + RNA	90.34	95.42	54.7	80.15
Probe 8 + TE Buffer	10.21	12.55	9.372	10.71
TE Buffer + RNA	5.579	6.509	6.388	6.159

These data show that a strong light signal is seen when the reaction mixes containing probes 6 or 8 and target RNA were added to the L/L reagent but little signal was seen when the probes were incubated without target RNA, or when the target RNA was incubated without these probes. In addition, probes 5 and 7 provided very low signals in the presence or absence of added target RNA. Probes 6 and 8 were designed to complement two different regions in the coding region of globin mRNA. Probes 5 and 7 were made to exactly mimic the sequence of these same target RNA regions. Thus, these data provide a second example of how the pyrophosphorylation of a probe can be used to detect a specific RNA.

Probe 5	SEQ ID NO:25	5'ATGGTGCATCTGTCCAGTGAGGAGAA GTCT3'
Probe 6	SEQ ID NO:26	5'AGACTTCTCCTCACTGGACAGATGCA CCAT3'
Probe 7	SEQ ID NO:27	5'GCTGCTGGTTGTCTACCCATGGACCC 3'
Probe 8	SEQ ID NO:28	5'GGGTCCATGGGTAGACAACCAGCAGC 3'

Example 7:        Specific Detection of RNA:  
Comparison of Signals from RNA Species  
that Match Probe Sequences in  
Reactions With and Without Added  
Extraneous Target RNA

For the pyrophosphorylation reaction  
described in Example 8 to be used to detect specific  
target sequences, another requirement of the system  
is that the probes should give a very similar signal  
in the presence and absence of extraneous RNA. In  
this example, the strength of the signal of probes  
designed to detect target globin mRNA in the presence  
of a large amount of yeast RNA is compared to the  
signal seen in the absence of added yeast RNA.  
Hybridization solutions containing various levels of  
yeast RNA, Probe 6 (SEQ ID NO:26) or Probe 8 (SEQ ID  
NO:28) and target globin mRNA (Gibco BRL, 18103-028)  
were assembled by adding 5  $\mu$ L 500 ng/ $\mu$ L either probe  
6 or probe 8 to 5 $\mu$ L 40 ng/ $\mu$ L of target globin mRNA  
and 10  $\mu$ L yeast RNA (Sigma Chemical Co. R3629) in 1X  
TE buffer (10 mM Tris, 1 mM EDTA) to produce  
solutions containing total amounts of yeast RNA of 0,  
2, 20, 200, 400, and 800 ng. The solutions were



heated at 50°C for 15 minutes and then permitted to cool to room temperature for 15.

Reaction master mix was assembled as in Example 2 above and 18 µL of the mix were placed in 18 tubes. After cooling 15 minutes, 2 µL of the various hybridization solutions containing probe 6 were added to the tubes and the tubes were placed in a 37°C heating block.

After 15 minutes of incubation of the hybridization mixture with the reaction master mix, 20 µL of the solution were added to 100 µL of L/L reagent (Promega, F202A) and the light output of the resulting reaction was measured using a Turner® TD-20/20 luminometer.

After the probe 6 data were collected, an identical set of reactions was performed using the hybridization solutions containing probe 8. The following data were obtained:

Probe 6 Reactions

Yeast RNA	relative light units			Average
None	96	109	111	105.3
2 ng	98.4	85.0	118.5	100.7
20 ng	117.9	110.9	82.7	103.65
200 ng	56.4	110.1	93.2	86.6
400 ng	115.7	110.7	124.6	117
800 ng	127.6	128.7	143.1	133.1

Probe 8 Reactions

Yeast RNA	relative light units			Average
None	105.8	97.0	82.3	95.0
2 ng	84.5	84.6	93.7	87.6
20 ng	99.6	111.7	104.9	105.4
200 ng	83.6	75.9	95.6	85.1
400 ng	94.7	97.2	81.9	91.2
800 ng	50.7	89.0	82.1	73.9

These data indicate that addition of very large amounts of yeast RNA to the hybridization reaction does not greatly lower the signal from hybridized probes for specific target RNA species.

Probe 6	SEQ ID NO:26	5'AGACTTCTCCTCACTGGACAGATGCACC AT3'
Probe 8	SEQ ID NO:28	5'GGGTCCATGGGTAGACAACCAGCAGC3'

#### Example 8: Initial Detection Limit For Plasmid Target DNA By Use Of Probe Pyrophosphorylation

In the previous two examples, plasmid target DNA was specifically detected using probes that hybridized to a target sequence in the DNA. In this example, a titration of target DNA is carried out in the pyrophosphorylation reaction to determine the level of DNA needed to obtain a signal from this reaction.

The *Sph* I cut target pKAN DNA (40,000 pg/ $\mu$ L) was serially diluted using nuclease-free water to obtain concentrations of 10,000, 2,500, 625, 156 and 39 pg/ $\mu$ L. Duplicate solutions containing 1  $\mu$ L each of these DNA target solutions, 1  $\mu$ L Probe 1 (SEQ ID NO:21) and 18  $\mu$ L nuclease-free water were assembled as were a pair of solutions containing 1  $\mu$ L Probe 1 and 19  $\mu$ L of nuclease-free water. All of these solutions were heated at 95°C for 3 minutes and then cooled for 10 minutes to room temperature to permit hybridization and form a treated sample. A 2X Master Mix was assembled as follows:

40  $\mu$ L 10X DNA Polymerase buffer (Promega, M195A)  
10  $\mu$ L 40 mM Sodium Pyrophosphate  
10  $\mu$ L (10 U/ $\mu$ L) Klenow exo minus DNA Polymerase (Promega, M128B)  
2  $\mu$ L NDPK at a concentration of 1 U/ $\mu$ L  
4  $\mu$ L 10  $\mu$ M ADP  
134  $\mu$ L nuclease-free water

The Master Mix components were mixed and 20  $\mu$ L 2X Master Mix were added to each of the solutions and incubated at 37°C for 20 minutes. A sample containing 4  $\mu$ L of the solution was then added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light produced by the reaction was immediately measured using a Turner® 20/20 luminometer. The following data were obtained.

Data Table		
Reaction	DNA Assayed*	Light Units
#1	4000 pg	168.4
#2	4000 pg	169.4
#3	1000 pg	57.7
#4	1000 pg	77.9
#5	250 pg	19.3
#6	250 pg	21.1
#7	62.5 pg	6.3
#8	62.5 pg	6.4
#9	15.6 pg	2.4
#10	15.6 pg	2.3
#11	3.9 pg	1.4
#12	3.9 pg	1.4
#13	0 pg	1.1
#14	0 pg	1.4

\*This number reflects that relative amount of DNA transferred to L/L solution.

- 5                    These data demonstrate that the detection limit for DNA by this reaction under these conditions is at least about 62.5 pg of DNA and is more likely about 15.6 pg of DNA or less.

Probe 1	SEQ ID NO:21	5'GCAACGCTACCTTTGCCATGTTTC3'
---------	--------------	------------------------------

Example 9: Detection Of  $\beta$ -galactosidase  
Target Sequences In Plasmids

In this example, two probes are used that complement each other exactly. One of the probes matches the sequence of the  $\beta$ -galactosidase gene exactly (sense orientation) and the other probe exactly matches the complementary strand (antisense orientation) of that gene. This example demonstrates that, whereas both probes can be used to detect the presence of the target  $\beta$ -galactosidase gene in plasmid DNA, the level of background signal given by reactions containing only probe DNA can be very different.

Probe 23 (SEQ ID NO:29) and Probe 24 (SEQ ID NO:30) were dissolved as described above to a concentration of 500 ng/ $\mu$ L and then diluted in nuclease-free water to 100 and 20 ng/ $\mu$ L. Plasmid pGEM7zf+ (Promega) was digested with *Sac* I (Promega) as the target and diluted to give a solution containing 20 ng of plasmid target DNA/ $\mu$ L of solution.

The following solutions were assembled:

Solution	Plasmid DNA ( $\mu$ L)	Probe, Concentration	H <sub>2</sub> O ( $\mu$ L)
#1	1	(none, 1 $\mu$ L of 1 X TE buffer added)	18
#2	0	1 $\mu$ L Probe 23, 500 ng/ $\mu$ L	19
#3	0	1 $\mu$ L Probe 23, 100 ng/ $\mu$ L	19
#4	0	1 $\mu$ L Probe 23, 20 ng/ $\mu$ L	19

Solution	Plasmid DNA (μL)	Probe, Concentration	H <sub>2</sub> O (μL)
#5	1	1 μL Probe 23, 500 ng/μL	18
#6	1	1 μL Probe 23, 100 ng/μL	18
#7	1	1 μL Probe 23, 20 ng/μL	18
#8	0	1 μL Probe 24, 500 ng/μL	19
#9	0	1 μL Probe 24, 100 ng/μL	19
#10	0	1 μL Probe 24, 20 ng/μL	19
#11	1	1 μL Probe 24, 500 ng/μL	18
#12	1	1 μL Probe 24, 100 ng/μL	18
#13	1	1 μL Probe 24, 20 ng/μL	18

These solutions were heated at 95°C for 3 minutes, and cooled to room temperature to form hybrids and treated samples. Then, 20 μL of 2X Master Mix made as described in Example 8 were added and the solutions incubated for another 20 minutes at 37°C. Four microliters of each solution were then added to 100 μL of L/L reagent (Promega, F202A) and the light output of the reaction immediately measured using a Turner® TD20/20 luminometer. The following data were obtained.

Reaction	Light Output	Net Light Output*
#1	2.8	
#2	4.0	
#3	1.9	
#4	1.3	
#5	52.4	45.6

Reaction	Light Output	Net Light Output*
#6	13.6	8.9
#7	4.1	0
#8	34.3	
#9	6.6	
#10	1.7	
#11	59.8	22.7
#12	19.3	9.9
#13	6.0	1.5

\*Net light output is calculated by subtracting the probe alone and DNA alone values from that obtained with both components present.

5

These data indicate that both probes can be used to generate a signal indicating the presence of the target region encoding the  $\beta$ -galactosidase gene matching the probes is present in the plasmid. They also demonstrate that the level of signal produced with a probe in the absence of target DNA can vary and that the signal from a probe and the complement of that probe are not necessarily equal.

10

\*

Probe 23	SEQ ID NO:29	5'CAGTCACGACGTTGTAAAACGACGGCC AGT3'
Probe 24	SEQ ID NO:30	5'ACTGGCCGTCGTTTTACAACGTCGTGA CTG3'

15

Example 10: Detection of Specific Target

DNA Sequences On Lambda DNA

In this example, detection of the target  $\beta$ -galactosidase gene in the DNA of a recombinant Lambda phage is demonstrated.

Duplicate solutions were made that contained: Solution 1 and 2, 1  $\mu$ L 300 ng/ $\mu$ L of Lambda gt11 DNA and 19  $\mu$ L of nuclease-free water; Solution 3 and 4, 1  $\mu$ L 500 ng/ $\mu$ L Probe 23 (SEQ ID NO:29) and 19  $\mu$ L nuclease-free water; Solution 5 and 6, 1  $\mu$ L 300 ng/ $\mu$ L Lambda gt11 DNA, 1  $\mu$ L 500 ng/ $\mu$ L Probe 23, and 18  $\mu$ L of nuclease-free water. All of these solutions were heated at 95°C for 3 minutes and then cooled to room temperature for 10 minutes to permit hybridization to occur between complementary strands and form treated samples. At this point, 20  $\mu$ L of 2X master mix made as described in above in this example were added and the solutions incubated for another 20 minutes at 37°C. A 4  $\mu$ L sample of each pyrophosphorolysis reaction was then taken and added to 100  $\mu$ L of L/L reagent (Promega, F202A) and the light production of the solution immediately measured with a Turner® TD20/20 luminometer. The following data were obtained.

Reaction	DNA Components	Light Units
#1	Target Lambda DNA	16.5
#2	Target Lambda DNA	7.4
#3	Probe 23	2.9



Reaction	DNA Components	Light Units
#4	Probe 23	2.9
#5	Target Lambda DNA and Probe 23	88.1
#6	Target Lambda DNA and Probe 23	70.4

These data indicate that the  
pyrophosphorylation system can be used to detect a  
probe hybridized to specific target sequences on  
5 lambda gt11 DNA.

Probe 23	SEQ ID NO:29	5' CAGTCACGACGTTGTAAAACGACGGCCA GT3'
----------	-----------------	---

10 Example 11: Detection of DNA Sequences in the  
Genome of *Campylobacter jejuni*

Oligonucleotides 11453 (SEQ ID NO:31) and  
11454 (SEQ ID NO:33) are exactly complementary and  
can be annealed, thereby forming a synthetic target  
15 representing a 70 bp segment of *Campylobacter jejuni*.  
These two oligonucleotides were diluted in nanopure  
water to a final concentration of 10 µg/mL. Four  
microliters of each were then mixed with 232 µL 10 mM  
Tris pH7.3 to yield a target solution of 0.3 µg/mL of  
20 DNA. Oligonucleotides 11451 (SEQ ID NO:32) and 11450  
(SEQ ID NO:34) are *Campylobacter jejuni* interrogation  
probes that bind to opposite strands of the bacterial